


Original article

Plasmid carriage and the natural complexity of bacterial populations contributes to plasmid persistence

 Tatiana Hillman ^{a,*} 
^a LAL4Bsynbiotics L.L.C., Biotechnology, Los Angeles, California, United States

ARTICLE INFO
Article history:

Received 23 February

2022

Received in revised form

02 June 2022

Accepted 11 June 2022

Keywords:

Plasmid carriage

Plasmid persistence

Biofilm

Conjugation

HGT

ABSTRACT

Introduction: Plasmids carry and transport genes that assist their hosts to survive in many environments. Many studies have examined the conditions for plasmid persistence in bacterial populations. A limitation includes that a majority of the mathematical models for examining plasmid persistence only included bacteria from similar colonies. However, most bacterial cells inhabit complex communities where plasmids disseminate between varied bacterial host cells. Thus, there is a gap in knowledge concerning the persistence of plasmids in natural bacterial populations. To address a few of these gaps in knowledge, the present study attempted to examine the effects of plasmid carriage on intrinsic stages of bacterial populations in *Bacillus subtilis* co-cultures.

Material and methods: *B. subtilis* cells were transformed with CRISPR-hCas-9 plasmid vectors where the natural phases of bacterial growth, biofilm production, and antibiotic resistance were examined in relation to plasmid carriage. These three natural phases were measured in relation to plasmid carriage through *in vitro* co-culture assays.

Results: After calculating the CFU/mL, bacterial growth in the *B. subtilis*-Carrier with *Escherichia coli* (B. sub-C-*E. coli*) and *Vibrio harveyi* (B. sub-C-VH) co-cultures significantly decreased with a paired-t-test two-tailed $P=0$. The WT *B. subtilis*-V.H samples, the *B. subtilis* Carrier-V.H co-cultures, and the controls each scored a total of 40, 47, and 46 of crystal violet (CV) intensity of biofilm, respectively. Biofilm formation decreased after co-culturing *E. coli* with the *B. subtilis*-Carrier, yielding a $P<0.001$. The antibiotic resistance levels of the co-cultures increased by 3% for the B. sub-C-V.H samples while the B. sub-C-*E. coli* co-cultures decreased in antibiotic sensitivity by approximately 1.5%.

Conclusions: Plasmid carriage contributes to plasmid persistence via altering the natural phases of bacterial populations.

© 2022 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author.

E-mail address: thillman@thel4binc.org

ISSN: 2695-5075 / © 2022 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.53986/ibjm.2022.0024>

El transporte de plásmidos y la complejidad natural de las poblaciones bacterianas contribuyen a la persistencia del plásmido

INFO. ARTÍCULO

Historia del artículo:
Recibido 23 Febrero
2022
Recibido en forma
revisada 02 Junio 2022
Aceptado 11 Junio 2022

Palabras clave:
Trasporte de plásmido
Persistencia de plásmido
Biofilm
Conjugación
HGT

RESUMEN

Introducción: Los plásmidos portan y transportan genes que ayudan a sus huéspedes a sobrevivir en muchos entornos. Muchos estudios han examinado las condiciones para la persistencia de plásmidos en poblaciones bacterianas. Una limitación incluye que la mayoría de los modelos matemáticos para examinar la persistencia de plásmidos solo incluyeron bacterias de colonias similares. Sin embargo, la mayoría de las células bacterianas habitan en comunidades complejas donde los plásmidos se diseminan entre diversas células huésped bacterianas. Por lo tanto, existe un vacío en el conocimiento sobre la persistencia de plásmidos en poblaciones bacterianas naturales. Para abordar algunas de estas lagunas en el conocimiento, el presente estudio intentó examinar los efectos del transporte de plásmidos en las etapas intrínsecas de las poblaciones bacterianas en cocultivos de *Bacillus subtilis*.

Material y métodos: Células de *B. subtilis* se transformaron con vectores plasmídicos CRISPR-hCas-9 donde se examinaron las fases naturales de crecimiento bacteriano, producción de biopelículas y resistencia a los antibióticos en relación con el transporte del plásmido. Estas tres fases naturales se midieron en relación con el transporte de plásmidos a través de ensayos de cocultivo in vitro.

Resultados: Después de calcular las UFC/mL, el crecimiento bacteriano en los cocultivos de *B. subtilis*-Carrier con *Escherichia coli* (*B. sub-C-E. coli*) y *Vibrio harveyi* (*B. sub-C-VH*) disminuyó significativamente con un *t*-test de dos colas $P=0$. Las muestras WT *B. subtilis*-V.H, los cocultivos *B. subtilis* Carrier-V.H y los controles obtuvieron cada uno un total de 40, 47 y 46 de intensidad de biopelícula cristal violeta (CV), respectivamente. La formación de biopelículas disminuyó después de cocultivar *E. coli* con *B. subtilis*-Carrier, lo que arrojó un $P<0,001$. Los niveles de resistencia a los antibióticos de los cocultivos aumentaron un 3 % para las muestras de *B. sub-C-V.H*, mientras que las muestras de *B. sub-C-E*. Los cocultivos de *E. coli* disminuyeron la sensibilidad a los antibióticos en aproximadamente un 1,5 %.

Conclusiones: El transporte de plásmidos contribuye a la persistencia de plásmidos mediante la alteración de las fases naturales de las poblaciones bacterianas.

© 2022 Los Autores. Publicado por Iberoamerican Journal of Medicine. Éste es un artículo en acceso abierto bajo licencia CC BY (<http://creativecommons.org/licenses/by/4.0/>).

HOW TO CITE THIS ARTICLE: Hillman T. Plasmid carriage and the natural complexity of bacterial populations contributes to plasmid persistence. Iberoam J Med. 2022;4(3):143-156. doi: 10.53986/ibjm.2022.0024.

1. INTRODUCTION

Plasmids are external genetic components that can become mobile. Plasmids can become transferred between bacterial cells via conjugation. Plasmids carry and transport genes that assist their hosts to survive in a plethora of environments [1]. The ability of plasmids to carry and transfer genes can initiate and complete the evolution and ecology of bacterial cells [1]. Because plasmids disseminate antibiotic resistance genes, these antibiotic resistance modes of action can continue to persist among medical pathogens for many additional decades [2]. These plasmids act as a type of genetic parasite [1]. Antibiotic resistance genes serve as benefits encoded in plasmids that can lead to virulence, high tolerance of heavy metals, and to higher rates of carbon metabolism [1]. Plasmids can also alter the environment of surrounding bacterial cells where beneficial enzymes or sources of nutrients can be produced

[1]. Although plasmids are found in bacterial populations, plasmids can form many changes of physiology in their bacterial hosts, which then reduces plasmid fitness [3]. The cost to fitness has created a quandary and a difficulty for interpreting how plasmids are conserved in a bacterial population for the long-term [4]. The challenges of understanding plasmid persistence has led to an unsolved riddle known as the “the plasmid paradox” [4].

Many hypotheses have been presented to attempt to solve this paradox. For example, it has been argued that bacterial evolution fuels plasmid persistence by lessening the fitness cost combined with plasmid carriage where an amplified rate of conjugation solidifies the persistence and survival of plasmids [5]. As a result, the persistence and containment of plasmids in bacterial populations over time presents to the current field a plasmid persistence puzzle or a plasmid paradox. In theory, plasmids carrying beneficial genes do not provide a solid explanation for the maintenance of plasmids [6]. Because of fitness costs, plasmids in theory

should dwindle while the beneficial genes integrate within the chromosomes of bacteria. As a result, many additional hypotheses have been generated to more succinctly explain the paradox of plasmid persistence. These hypotheses include a higher frequency of HGT and conjugation during infectious transmission, the plasmid transport between strains of bacteria, and in the maintenance of the higher proliferating bacterial strains, which display amplified effectiveness of transconjugation [6].

For many decades, a large number of studies have examined the conditions for plasmid persistence and maintenance in bacterial populations. A limitation takes into consideration that a majority of the mathematical models for examining plasmid population biology research involves bacterial populations within similar colonies. However, most bacterial cells inhabit complex communities where plasmids disseminate through conjugation from varied bacterial hosts [1]. Therefore, there is a gap in knowledge concerning the persistence of plasmids in natural bacterial populations and communities [1-2]. It is crucial to resolve these limitations to enhance our understanding of plasmid persistence. To address a few of these gaps in knowledge, the present study attempted to examine plasmid persistence in relation to plasmid carriage in *Bacillus subtilis* co-cultures. *B. subtilis* inhabits soils and dwells in many environments and ecosystems. *B. subtilis* are also gut commensal bacteria found in animals and in humans [7]. *B. subtilis* was selected for this study because it uses plasmids to assist with HGT in the gut microbiome [7]. Moreover, more frequent HGT events occur in the human gut microbiome than in a diverse ecosystem of non-human microbes [8]. Physical proximity contributes the most to the large operation of HGT in the gut via cross-domain gene transfer [9]. Conjugation as a mechanism of HGT is substantially altered by physical proximity [9]. Conjugation is effective in the gut because it conserves and protects DNA from enzymes known as nucleases along with elimination of heavy metals in the environment [9]. Conjugation via HGT has a highly significant role in the gut microbiome, and an improved understanding of conjugation's contribution to HGT in the gut is greatly needed [9].

Therefore, the main objective of this study was to provide additional evidence and explanations for plasmid persistence. To offer a potential interpretation of plasmid persistence, a possible relationship between natural bacterial conditions and plasmid persistence were investigated in *B. subtilis* (B. sub), *Vibrio harveyi* (VH), and *Escherichia coli* (*E. coli*) bacterial cell co-cultures. These bacterial strains were selected to model *in vitro* and

naturally occurring communities of bacteria that represent an interaction between a few bacterial phylotypes present in the gut microbiome. For example, in this study, *B. subtilis* and *E. coli* that inhabit the gut microbiomes of animals and humans were used while *V. harveyi* was examined, which is a pathogen found in fish, in which many humans and animals consume. Factors and conditions, such as bacterial growth, biofilm formation, and antibiotic resistance genes, which may contribute to plasmid persistence, were analyzed in this study. These natural phases of bacterial populations were observed for any changes in bacterial host fitness as related to plasmid carriage that may offer insights into the dynamics and contributors of plasmid persistence. It was hypothesized that plasmid carriage can alter the natural phases of bacterial communities, which all interrelate with the plasmid persistence of bacterial host cells. To test the hypothesis of this study, *B. subtilis* cells were transformed with Clustered Regularly Interspaced Short Palindromic Repeats or CRISPR plasmid vectors encoding human CRISPR associated protein 9 (hCas9) of *accA*, which codes for the bacterial acetyl-CoA carboxylase subunit *AccA*. The plasmid utilized a promoter exclusive only for human RNA polymerase to bind so that the bacterial *accA*-sgRNA could not be expressed. This was done in order to only observe the effects of plasmid carriage without gene modifications in the target bacterial host. After transformation of *B. subtilis* with the CRISPR-hCas9-*accA* plasmid vectors, the *B. subtilis* plasmid carriers and non-carriers (B.sub-C/B.sub-NC) were co-cultured with *E. coli* and *V. harveyi* (VH) in order to exhibit any changes in the conditions of the bacterial cell co-cultures. The bacterial cell growths, biofilm intensity, and the antibiotic resistance of the co-cultures were then examined via common microbiological assays such as a biofilm crystal violet assay. As a consequence, the purpose for this study was to observe and analyze possible significant changes in the innate phases of bacterial communities to find a significant link between the three aforementioned natural bacterial states and the plasmid carriage of the bacterial co-cultures, which all can maintain plasmid persistence.

2. MATERIAL AND METHODS

2.1. BACTERIAL STRAINS

Bacterial strains included *B. subtilis*, *V. harveyi* DSM 6904 (DSMZ), *E. coli* BL21, and *E. coli* DH5a. Luria Bertani broth and agar were inoculated with each strain of bacteria aforementioned. The LB liquid and LB agar media were

incubated from 16 hours to 24 hours at the temperatures specific for each bacterium. *B. subtilis* and *V. harveyi* bacterial cells were incubated at 30 degrees Celsius as *E. coli* cell cultures were grown at 35 to 37°C.

2.2. PLASMID DESIGN AND ASSEMBLY

The vector named pRP[CRISPR]-hCas9-U6>{accA} was designed in VectorBuilder.com. The vector ID is VB210702-1210yff with a total value of 8431 base pairs (bp). The vector was designed to contain a single guide RNA (gRNA) for CRISPR-hCas9 to carry the 20bp of the *accA* gene. The gRNA can express the sequence of CTGCGTGAAATGTCTCGCCT, which is a guide RNA of 20bp for the bacterial *accA* gene. To assemble pRP[CRISPR]-hCas9-U6>{accA} hCas9 was inserted into the vector. The antibiotic resistant gene for ampicillin was also included in the plasmid assembly. The plasmids were produced at a high plasmid copy number. The fully assembled vectors were assembled and delivered by VectorBuilder. The plasmids were delivered through a glycerol stock solution of *E. coli* cells. The pRP[CRISPR]-hCas9-U6>{accA} plasmids were purified from the *E. coli* cells via the BioBasic EZ-10 Spin Column plasmid DNA Kit (BS413-50 preps).

2.3. B. SUBTILIS TRANSFORMATION

The pRP[CRISPR]-hCas9-U6>{accA} plasmids isolated from the *E. coli* stock of cells were used to transform *B. subtilis* cells, deriving the *B. subtilis*-Carriers. To optimize the *B. subtilis* transformation protocol, *B. subtilis* was cultured for 5 hours until the exponential phase of growth with an OD600 value of 0.3-0.4. Two millimeters were taken from the culture and centrifuged for 1 minute at 8,000 rpm, collecting the *B. subtilis* cells. The supernatant was discarded. About 200 µL of LB broth with 2% xylose was added and then vortexed to concentrate the cell suspension at 10x. The 2% xylose was added to the LB. The isolated pRP[CRISPR]-hCas9-U6>{accA} plasmids with ampicillin selective markers were added to the concentrated *B. subtilis* cells with the 2% xylose-LB solution. The *B. subtilis* transformations were incubated for 1 hour at 37°C and then plated. About 20 µL each of the pRP[CRISPR]-hCas9-U6>{accA} *B. subtilis* transformations were inoculated onto two LB agar plates and 160 µL were also plated. The plates were sealed with parafilm and incubated overnight at 37°C. Each LB agar culture plate contained carbenicillin to select for the pRP[CRISPR]-hCas9-U6>{accA} plasmids with the ampicillin selective marker because carbenicillin cannot degrade in high acidity or in excessive heat.

2.4. CO-CULTURE ASSAYS

The *V. harveyi* and the *E. coli* bacterial cells were co-cultured with the *B. subtilis* Carriers and the wild type (WT) *B. subtilis* to detect any change in growth or bacterial host fitness of the *B. subtilis*-Carriers. After completing the co-cultures, the *B. subtilis*-Carrier co-cultures with *V. Harveyi* and *E. coli* BL21/DH5α cells were compared to the co-cultures of the WT *B. subtilis* with the *V. harveyi* and the *E. coli* BL21/DH5α. To use a co-culture assay and approach, each bacterial species and strain, aforementioned, were subcultured on separate agar plates. Colonies from the agar plates were inoculated into LB liquid medium and then incubated overnight. About 5-6 bacterial colonies from each *E. coli* BL21/DH5α and *V. harveyi* plate and 5-6 colonies from the *B. subtilis*-Carrier and WT *B. subtilis* agar plates were inoculated into LB broth. The co-cultures without the *B. subtilis* carriers were used as the control samples. The co-cultures were grown overnight at 30 to 35°C and then analyzed for CFU/mL, biofilm production, and antibiotic resistance.

2.5. COLONY FORMING UNITS (CFU)

To measure the total of colony forming units (CFU), 10µL of each stock of the bacterial strains were pipetted into 1.5 mL Eppendorf centrifuge tubes. About 0.1mL of the 5 mL stock was added to 0.09 mL of LB broth to form the 10⁻¹ dilution of each bacterial strain. The dilutions were resuspended. Then, 10µL of the 10⁻¹ dilution was added to 90µL of LB broth to form a dilution of 10⁻². The 10µL of 10⁻² were added to 90µL of LB broth, and then more dilutions were continued in a series to a total of 10⁻¹⁰ dilutions. The last dilution and two other dilution series were selected for plating. The plates with the selected dilutions were incubated overnight at 37°C. The number of colonies for each bacterial strain was counted from the cultured agar plates, using the Promega Colony Counter computer software program. The number of colonies counted was substituted into an equation for calculating the values of CFU for bacterial cells.

The equation included:

$$\text{CFU/mL} = [\text{Number of Colonies Counted}] \times [\text{Dilution Factor}] / \text{Volume spread on plate}$$

2.6. KIRBY-BAUER DISK DIFFUSION TESTS

The co-cultures of *E. coli* BL21/DH5α and *V. harveyi* with *B. subtilis* Carriers and WT *B. subtilis* were then assayed for antibiotic resistance. The Kirby-Bauer Disk Diffusion Tests were used. Carbinicillin, chloramphenicol,

tetracycline, and kanamycin were pipetted into sterilized antibiotic disks. Approximately, 5 μ L of each of the four antibiotics were pipetted into the antibiotic disks. A lawn of each bacterial strain was inoculated on LB agar plates via the spread plate method. A swab of LB broth liquid cultures from each bacterial strain was spread onto agar culturing media. The antibiotic disks were placed onto the bacterial culture plates. The plates were incubated for 24 hrs at 35°C. The zones of inhibition were measured in millimeters (mm) and then the values were interpreted, using a zone diameter interpretive standards chart provided by Sarker et al. [10].

2.7. BIOFILM ASSAY

Five milliliters of LB broth cultures were prepared from each bacterial strain and co-culture. The biofilms of the *E. coli* and *V. harveyi* strains were used as the controls and were co-cultured with the *B. subtilis* Carriers and WT *B. subtilis*. The cultures grew for 20 hours at 37°C. The 1: 100 dilution sets were prepared, totalling to 1mL of the LB/Carb liquid cultures. About 100 μ L of the dilution series were pipetted into 4 wells, per bacterial strain, of a 96 well plate. The plates were incubated for 48 hours at 37°C. The plates were shaken to remove any planktonic bacteria. The plates were rinsed with water. All the wells were stained with 125 μ L of 0.1% crystal violet solution for 10 min. The 96-well plate was shaken and rinsed with water to remove any excess crystal violet solution.

2.8. STATISTICAL ANALYSIS

The p-Values of Colony Forming Units for *V. harveyi* samples were analyzed through a Paired-Two-Tailed t-test. The biofilm assays were analyzed via Two-Way ANOVA. The antibiotic test data results were analyzed by One-Way ANOVA for *E. coli* samples as paired-t-tests, which were used to generate p-Values for *V. harveyi* antibiotic resistance sample results. Descriptive and inferential statistics were performed, using Microsoft Excel and GraphPad Prism. All graphs and other statistical analyses were generated through ChartExpo and GraphPad Prism.

3. RESULTS

3.1. BACTERIAL GROWTH DECREASED IN *V. HARVEYI*, *E. COLI*, AND *B. SUBTILIS*-CARRIER CO-CULTURES

After transforming *B. subtilis* cells with the CRISPR-hcas9-sgaccA plasmid vectors, the transformants were co-

cultured with *V. harveyi* and *E. coli*. The *B. subtilis* cells that were transformed with CRISPR-hCas-9 plasmid vectors were compared against bacterial samples with non-carriers such as the *V. harveyi* and the *E. coli*. The colony forming unit (CFU) values of the non-carriers were compared to the carriers. As a result, after calculating the CFU/mL, bacterial growth, in the *B. subtilis*-Carrier and *V. harveyi* co-cultures, significantly decreased with a paired-t-test two-tailed $P=0$ (Figure 1A). *V. harveyi* decreased by an average of 31% after the co-culture assays (Figure 1A). The bacterial growth between *E. coli* and *B. subtilis* Carrier co-cultures produced a percent difference of 54% (95% CI [3.06M, 1.10M]). *E. coli* yielded 3.02E7 CFU as the co-culture decreased to 1.10E7 CFU. *V. harveyi* had an average CFU of 5.8E11 that was then reduced to 3.4E11 CFU during the co-culture assays (Figure 1A). Overall, the co-cultures containing *B. subtilis* plasmid carriers exhibited lesser bacterial growth (Figure 1B).




Figure 1: The Colony Forming Units (CFU/mL) After dilution series for each sample were generated, the CFU/mL were allocated. Each dilution series showed a large decrease in CFU/mL for *B. subtilis*-Carrier-*V. harveyi* compared against the *V. harveyi* only bacterial sample culture. A. A significant decrease in bacterial growth occurred for the *B. subtilis*-Carrier-*V. harveyi*-*E. coli* co-culture assays. B. *V. harveyi* CFU/mL dwindled from 3.69B to 2.46B when *V. harveyi* was co-cultured with *B. subtilis* plasmid carriers. *E. coli*, when co-cultivated with plasmid carrier *B. subtilis*, displayed a significant decrease in CFU/mL from 3.02M to 1.10M.

3.2. BIOFILM FORMATION WAS AMPLIFIED IN *B. SUBTILIS* CARRIER-VH CO-CULTURES, BUT REDUCED IN *E. COLI* CO-CULTURES

Biofilms include populations of microorganisms growing on a surface within a matrix of extracellular polymeric molecules. An innate property of biofilms is that biofilms are spatial and organized shelters where microbes interact with adjacent microbial communities rather than with wider spaced bacteria. The spatial structural complex offers heterogeneity within these spatial structures of biofilm where bacteria experience different environmental conditions and pressures in different locations of the biofilm [11]. The spatial structure and the heterogeneity of this biofilm habitat significantly affect the evolution of populations of microbes [11]. For example, through slowing down competition between bacterial cells, biofilm spatial structures' environments select for many favorable mutations and then delay the rate of adaptation [11]. Thus, the biofilm intensity of each sample, including WT--VH-E.coli, B.sub-C-VH-E.coli, and WTB.Sub-NC-VH-E.coli, was measured through a crystal violet assay. Biofilm formation was amplified in *B. subtilis* Carrier-V.H co-cultures, but reduced in the *E. coli* co-cultures. Biofilm is vital for the survival, persistence, and for the proliferation of bacterial cells. Biofilm serves to insulate and protect bacterial cells from external antimicrobials such as antibiotics and other antagonistic microbes. Biofilm is also a major contributor to the increased virulence and proliferation of pathogenic bacteria, and it also invigorates antibiotic resistance. The biofilm production of *V. harveyi* and *E. coli* co-cultures with wild type *B. subtilis* were compared to the *B. subtilis* plasmid carriers, *V. harveyi*, and *E. coli* co-cultures. The wild type bacterial strains of *V. harveyi* and *E. coli* were used as control samples for comparative analysis. For *V. harveyi* samples grown with *B. subtilis* Carriers, biofilm formation exceeded the levels of biofilm generated by *V. harveyi* combined with WT *B. subtilis*. A Likert scale ranging from 1 to 5 was used to quantify the intensity of crystal violet absorbance added to the samples. A score of 4-5 equaled a high intensity of crystal violet coloration, and a value of 3 to 4 represented a moderate intensity of crystal violet absorbance. If a sampling well did not appear with much coloration or no stain was formed, these samples were given a score of 0 to 1.

The *B. subtilis*-C-V.H co-cultures averaged a score of 3.9, and the WT *B. subtilis*-V.H samples averaged a 3.3 crystal violet (CV) assay score. The control of the WT *V. harveyi* averaged a 3.8 CV intensity score. WT *B. subtilis*-V.H samples, the *B. subtilis* Carrier-V.H co-cultures, and the




Figure 2: Biofilm Assay. A. The crystal violet stain of *B. subtilis*-Carrier-V.H appeared with the highest intensity. *B. subtilis*-Carrier-V.H and V.H showed a higher score of CV intensity than the WT-*B. subtilis*-V.H co-cultures. B. The biofilm assays were performed in triplicates, in which the *B. subtilis*-Carrier-V.H had the highest expression of CV absorbance intensity. *B. subtilis*-Carrier-V.H co-cultures showed a higher intensity of Crystal Violet absorbance versus the wild type *B. subtilis* as shown in the simulation of data and in the scatter plot.

controls each scored a total of 40, 47, and 46 of CV intensity, respectively (Figure 2A-2B). The p-Values, after performing a Two-Way ANOVA analysis of each group's CV scores, computed to $P=0.134$, $P<.001$, and $P=0.64$ for comparing the difference between the scores, the difference between the bacterial sampling groups, and for determining the variable difference between the bacterial groups and their CV scores, respectively. There was an increase in biofilm yield when comparing the *B. subtilis*-C-V.H treatment groups against the WT *B. subtilis*-V.H sample groups. However, the CV scores, for the *B. subtilis*-C-V.H co-cultures, of biofilm formed were equally approximate to the WT-V.H sample scores. Biofilm formation decreased after co-culturing *E. coli* with *B. subtilis*-Carrier, yielding a $P<.001$ (Figure 3A-3C).




Figure 3: Biofilm Assay for *E. Coli* and *AB. Subtilis* Carrier A. *E. Coli* had a total score of 16 with an average CV intensity of 4 (Avg=4, SD=0). B. The Pearson r analysis shows coefficients of 1 in the non-carriers while there was a coefficient of 1 in the co-culture samples of carriers. There was a 0.71 coefficient between the noncarriers and the carriers, showing a decrease in correlations between each sample. C. *B.subtilis*-Carriers combined with *E. Coli* averaged a score of 1 CV intensity and a total score of 4 (Avg.=1 SD=0.8). *E. Coli*, grown with WT *B. subtilis*, averaged a 3.5 CV score and had a total score of 14 (Avg=3.5, SD=0.58).

3.3. B. SUBTILIS CARRIER-V.H AND B. SUBTILIS CARRIER-E. COLI CO-CULTURES SHOWED MORE ANTIBIOTIC RESISTANCE THAN CONTROL SAMPLES

The Kirby-Bauer antibiotic disk assay was used to analyze the antibiotic resistance of *B. subtilis*-Carrier co-cultures of *B.sub-C-VH-E.coli* and the non-carriers such as the *B.sub-NC-VH--E.coli*, including the control samples of WT-VH and WT-*E.coli*. The antibiotic resistance of *B. subtilis*-Carrier-V.H and the *V. harveyi* cultures were closely identical (Figure 4A). In fact, the *B. sub-C-V.H* cultures seemed to share the exact amount of antibiotic resistance with nearly equal zones of inhibition in millimeters. All co-cultures were resistant to carbenicillin and moderately resistant to tetracycline (Figure 4A). The *B. sub-C-V.H*, the WT-V.H, and the control samples were all sensitive to chloramphenicol and to kanamycin. The antibiotic resistance of *B. sub-C-V.H* co-cultures compared against the control samples had a Two-Tailed $P=0.51$ after using a paired t-test analysis. The *E. coli* samples showed similar levels of antibiotic resistance as the *B. sub-C-E. coli* co-cultures. All *B. sub-C-E. coli* sample groups were more

resistant to each antibiotic. The *B. subtilis-C-E.Coli* compared against *E. Coli*, after a One-Way ANOVA analysis, yielded a $P=0.49$. The antibiotic resistance levels of the co-cultures seemed to mimic the antibiotic resistance expressed by the control sample groups, which included *V. harveyi* or *E. Coli*; however, the antibiotic resistance increased by 3% for the *B. sub-C-V.H* samples while the *B. sub-C-E. coli* co-cultures decreased in antibiotic sensitivity by approximately 1.5% (Figure 4B). The *B. subtilis* carriers co-cultured with *V. harveyi* and *E. coli* displayed a higher level of antibiotic resistance than the controls and its wild-type because the *B. subtilis* carriers of the antibiotic resistant genes may have outcompeted the bacteria without the resistant plasmids (Figure 5).




Figure 4: Antibiotic Resistance Tests A. The top plates include the antibiotic resistant test results for *E. Coli* co-cultures and the bottom plates represent the results for *V. harveyi* sample groups. The top plates showed more moderate sensitivity to resistance of the antibiotics carbenicillin, chloramphenicol, tetracycline, and kanamycin. The WT *E. Coli* was more sensitive to tetracycline, chloramphenicol, and to kanamycin than the WT *B. subtilis* and the *B.subtilis*-Carrier-*E. Coli* co-culture. *V. harveyi* on the bottom plates showed identical and similar antibiotic resistance and sensitivity as the *B.subtilis*-Carrier-V.H and WT *B. subtilis* co-cultures. B. The zones of inhibition were measured in millimeters. The zone of inhibition for *V. harveyi* totaled 99 and 91 for *B. subtilis*-Carrier-V.H producing a 3.0% decrease in the zones of inhibition between each of these samples. The WT *E. Coli* had a total zone of inhibition of 8.5% that was raised to 10.1% for the *B. subtilis*-Carrier-*E. Coli* that yielded a total Zol of 33. The wild-type *V. harveyi* and *E. Coli* samples showed increased antibiotic sensitivity, however, the *B. subtilis*-Carrier-V.H and *B. subtilis*-Carrier-*E. Coli* co-cultures increased in antibiotic resistance.




Figure 5: Zones of Inhibition (mm) The zones of inhibition were measured in millimeters. The zone of inhibition for *V. harveyi* totaled to 99 and 91 for *B. subtilis*-Carrier-V.H. WT *E. Coli* had a total zone of inhibition of 38 as *B. subtilis*-Carrier-*E. Coli* yielded a total Zol of 33. Wild-type *V. harveyi* and *E. Coli* samples showed increased antibiotic sensitivity, however, the *B. subtilis* Carrier-V.H and *B. subtilis* Carrier-*E. Coli* co-cultures increased in antibiotic resistance.

4. DISCUSSION

4.1. PLASMID CARRIAGE, BACTERIAL GROWTH, AND FITNESS

Many studies have shown that for plasmid persistence to occur there must be an association between plasmid loss, transfer through conjugation, and the effects of plasmid carriage on the fitness of the host. Gama et al., concluded that regardless of the multiple types of interactions affecting the persistence of plasmids, there is a more potent effect on plasmid maintenance than on those interactions impacting conjugation and plasmid loss [12-17]. Additionally, Krone et al. used simulations of bacterial colony growth rates of plasmid carriers and plasmid-free strains as approximations for the growth rate parameters in their mathematical model [18]. They found a linear association between the radii of plasmid-carriers, plasmid-free bacteria, and cell growth rates [18]. As a result, in this study, the *B. subtilis* plasmid carrier co-cultures showed lesser levels of bacterial cell growth because plasmid carriage many times reduces the fitness of its host [5]. The frequency and rates of these dynamics depend on the plasmid, the environmental pressures, and on the host. Plasmid loss during bacterial growth can become sensed and detected [5]. For these reasons, the B.sub-C-VH and B.sub-C-*E. coli* co-cultures may have had lesser bacterial growth because of having a higher fitness cost. However, without plasmid loss or lacking any fitness cost, plasmids would still become removed from the populations of

bacteria [5]. Plasmid persistence, thus, depends more on fitness cost and horizontal transfer via conjugation [5].

The fitness that is related to plasmid carriage immensely affects the plasmid persistence in bacterial populations [1-5]. However, the effects of plasmid fitness on wild-type bacterial host cells have not been completely surveyed [1-5]. Computational experiments have shown that fitness varies between bacterial strains in a community, which then increases the probability of plasmid persistence at low rates of conjugation [1-5]. For these reasons, the co-cultures with plasmid carriers may have shown higher fitness costs through displaying a bacterial growth reduction. In this study, the frequency of plasmid transfer may have been reduced as the amount of diverse bacterial strains in the cultures increased because the reception of plasmids was linked to a cost of fitness [1-5]. In addition, the acquisition of plasmids can be amplified when the allocation of fitness effects becomes more complex with a wider range of variance [1-5]. A possible interpretation of these effects is that the cost of plasmid fitness was similar between all bacterial strains in the cultures without plasmid carriers where higher rates of plasmid persistence occurred. The higher rates of plasmid persistence were observed in the cultures without carriers because there were lower rates of conjugation combined with lower fitness cost compared to the bacterial cultures with the more diverse bacterial strains, which required higher rates of conjugation [1-5]. For example, in the Benz et al. study they represented their model for the spread and proliferation of plasmids in a complex environment such as in the gut of animals and humans that predicted the rate of transfer using *in vitro* testing and genetic data analysis [19]. Benz et al. removed ESBL-plasmids from their native hosts, and they confirmed that the donor and recipient strains worked to regulate the frequencies of conjugation transfers [19]. However, the rate of conjugation needed for plasmid persistence may be much lower than confirmed in past studies [1-5]. An example includes a mathematical model that was created for the fitness effects of a carbapenem-resistance plasmid in 50 wild-type enterobacteria found in the gut microbiota of humans [1-5]. Alonso-Valle's mathematical model showed that the chance of plasmid persistence is not dependent on an escalated rate of conjugation. This effect yields plasmids at a low average cost, but forms incomplete plasmids at a larger cost of fitness [1-5]. The chance for plasmid persistence amplifies with the increased numbers of bacterial strains, creating a highly complex community [1-5]. As the community increases in complexity, the stability of plasmid proliferation also elevates through the transfer dynamics of the plasmids [1-5]. This in turn offers

many available plasmid-permissive bacterial phylotypes [1-5].

In contrast, when the fitness effects differ, more available bacterial host cells in a population amplifies the chance of the uptake of plasmids by these host cells where no costs or favorable fitness effects are produced [1-5]. As a consequence, the *B. sub-C-VH* and the *B. sub-C-E. coli* co-cultures in this study may have decreased the availability of plasmid-free cells and exhibited reduced growth rates because of a greater exertion of a host fitness cost compared to the control samples. This evidence is significant because it confirms that plasmid carriage may have lessened the numbers of available plasmid-free bacterial cells needed for conserving host cell fitness and maintaining plasmid persistence. For instance, when plasmids are readily available in a complex bacterial community, plasmid persistence may continue even during the lack of conjugation [1-5]. Therefore, increasing the complexity of bacterial communities can increase the probability and continuance of plasmid persistence in natural environments [1-5]. Because many natural microbiomes consist of complex bacterial strains and communities, plasmids can easily proliferate within these natural environments [1]. Consequently, in this study, plasmid persistence may have occurred more frequently because of increased complexity present in the *B. subtilis* Carriers co-cultured with additional bacterial strains of *V. harveyi* and *E. coli*.

4.2. PLASMID CARRIAGE AND BIOFILM

More biofilm may have been produced in *B. sub-C-VH* treatment samples than in the *B. sub-C-E. coli* samples because of a greater presence of conjugative plasmids in the *B. sub-C-VH* samples than in the *B. sub-C-E. coli* cultures because conjugative plasmids improve the bacterial capability to form biofilms by using greater numbers of pili, which increases the adhesion within a biofilm [20]. Biofilm formation may have been higher in the *B. sub-C-VH* co-cultures than in the *B. sub-C-E. coli* because of the rate of conjugation that created more pilli formation. This increased pilli activity may have increased the adhesion in the biofilms of co-cultures with plasmid carriers. However, plasmids may still interplay with each other by organizing or inhibiting the transfer of plasmids [20]. Gama et al. examined and quantified the formation of biofilm in *E. coli* cells carrying two types of plasmids [20]. Gama et al. used 11 innate and extracted conjugative plasmids where they confirmed that a few plasmids promote the formation of biofilm through conjugation efficiency [20]. Gama et al. also confirmed the effect of

paired plasmids on biofilm formation. Their study showed increased formation of biofilm when both plasmids inhabited the same bacterial cell or when the plasmids were contained and carried in different cells [20]. One of the plasmids showed dominance over the other plasmid where one of the plasmids directed the increase of biofilm formation. *E. coli* cells showed low formation of biofilm in a crystal violet biofilm assay versus in quinolone-resistance *E. coli* found in the same population [21]. Acquiring the two different plasmids of the *E. coli* QREC virulent strain lowered the production of biofilm in the microtiter plates [21]. The rates of motility were reduced, but planktonic growth continued to increase [21]. Therefore, Nesse et al. hypothesized that plasmids carrying antibiotic resistance genes may have reduced the formation of biofilm dependent upon the fluxes of flagellar protein expression [21]. Likewise, in this study, the co-cultures of *E. coli* combined with *B. subtilis* plasmid carriers, containing the resistance gene of ampicillin, showed less biofilm generation possibly due to a lesser production of flagellar protein expression from the *E. coli* cells [21]. Currently, methods are needed in future research studies to better interpret the molecular basis of these dynamic plasmid interactions [20].

Nesse et al. also observed that the acquisition of plasmids reduced the rate of motility in the transconjugants of the bacteria [21]. Earlier studies have confirmed this when the motility of *E. coli* strains increased after removing cephalosporin-resistance plasmids from the bacterial strains of *E. coli* [21]. The plasmids may have reduced the motility where the plasmids affected biofilm formation in the transconjugant bacterial cells by lowering the rate of motility [21]. These observations are supported by plasmids being acquired that did not affect the biofilm production on Congo Red (CR) agar plates, showing that the control of biofilm production depended on environmental conditions, which differed between the assays for biofilm [21]. The walls of the wells in a microtiter plate, at the liquid to air interface, of bacterial cells may have needed increased motility to reach the location of the biofilm [21]. This is opposite from the agar surface with biofilm formation because on agar surfaces intense motility levels may not be required for biofilm formation [21]. This is supported by previous studies of *Salmonella* flagellar mutants, in which these bacteria showed deformities in early biofilm creation in microtiter plates used for crystal violet biofilm tests [21]. In a subsequent study *Salmonella* displayed flagella that were not required for producing biofilm on CR-agar plates, but the lack of flagella did negatively affect the biofilm formed

on air to liquid, surface to liquid, or cell to cell liquid interfaces [21]. Therefore, in this study, the *E. coli* cells co-cultured with *B. subtilis* carriers may have not shown profound biofilm formation because its plasmid carriage, containing ampicillin resistance genes, lowered the rate of motility needed for forming biofilm in the liquid cultures of the microtiter plates during earlier stages. This observation was depicted in the crystal violet biofilm assays of the B.sub-C-E Coli plasmid carriers, in which the B.sub-C-E. *coli* samples exhibited less biofilm formation possibly due to less motility negatively affecting the movement of the bacterial cells in the liquid to air interface during the early stage of biofilm generation.

However, in vibrios, biofilm creation is more robust because vibrios are highly motile during the early stages of biofilm formation in viscous liquid [22]. Although *V. harveyi* was co-cultured with the *B. subtilis* plasmid carriers, in this study, the B.sub-C-VH carrier cultures continued to show a greater intensity of biofilm formation because the *V. harveyi* produced a more vigorous motility. Many bacteria use their flagellum to produce much motility in the beginning stages of forming biofilm by increasing the movement into the biofilm or on its surface, but in vibrios motility involves more than initial attachment [22]. For example, it was shown in a previous study that *Vibrio parahaemolyticus* utilizes its motor to decide rather to move on a surface or swim in extremely viscous types of media [22]. *V. parahaemolyticus* uses its polar flagellum to initiate biofilm formation, so motility is central for biofilm formation in vibrios as observed in previous biofilm models [22]. The two flagellar systems power flagella through swimming in liquid media while the lateral flagellum drives the swarming on solid surfaces [23-28]. The biosynthesis of the flagellar complexes is needed for swarm motility [29]. Bacterial cells from swarming colonies have more numbers of flagella than cells isolated from liquid cultures [29]. Wild-type *B. subtilis* are able to form biofilm complex structures on the surfaces of standing cultures of liquid [29]. However, researchers viewed deformities in these biofilm structures in *B. subtilis* mutants that lacked surfactin production [29]. *B. subtilis* produces surfactin as a self-defense mechanism. Bindel et al. found that because of a loss of swarming motility, their mutant *B. subtilis* sample of A164Δ10 could not form pellicles on their liquid cultures, but when they added proteases to the medium of A164Δ10 cultures, pellicles were produced [29]. The Bindel et al. study confirmed that there is a strong correlation between swarming motility and biofilm production [29]. Additionally, in this study, it may be inferred that the *B. subtilis* carriers and non-carriers co-

cultured with *E. coli* and *V. harveyi* may have lacked biofilm formation because there may have been less surfactin production. In the tested samples of B.sub-C-E.coli and WT-B.sub-NC-VH, lesser amounts of surfactin may have led to a decreased rate of the *B.subtilis* ability to swarm, which then reduced the intensity of biofilm formed in the liquid culture crystal violet assays. Less biofilm was produced by the B.sub-C-E. *coli* growths due to the propensity of *E. coli* to have decreased rates of motility in liquid-to-air standing cultures while vibrios like *V. harveyi* use more vigorous flagellar motility in highly viscous medium not as utilized in liquid standing cultures. This reduction in swimming motility combined with a lesser presence and adhesion of *E. coli* cells or *V. harveyi* cells in the samples may have negatively affected the surfactin production of the *B. subtilis* plasmid carriers and of the wild-type *B. subtilis* non-carriers in these subsequent co-cultures, leading to the observed decrease of biofilm formation. Roder et al. showed that biofilms are spatial storages for plasmids, which allow plasmids to persist even in intense pressures of environmental conditions [30]. Bacterial cells that carry plasmids increase the removal of bacterial cells lacking plasmids where biofilms can become temporary storages for the plasmids [30]. However, plasmid maintenance expends much energy and leads to a fitness cost of the host cell [30].

4.3. PLASMID CARRIAGE AND ANTIBIOTIC RESISTANCE

Antibiotic resistance increased in the *B.subtilis* carrier co-cultures because plasmids can be transferred when their rate of conjugation and horizontal transfer overrides the costs of plasmid carriage in a bacterial cell [31]. In a previous study, Svara et al. constructed a mathematical model to study the dynamics of plasmids as they responded to varied treatments of antibiotics [31]. However, when there were reduced fitness costs of plasmids, a beneficial transfer of plasmids dwindled [31]. Svara et al. produced a model for the absence of non-resistant plasmids as plasmids remain in a bacterial population when the rate of transfer is most efficient and favourable [31]. Svara et al. found that the resistant plasmids, such as the plasmids with antibiotic resistance genes, dominated the non-resistant plasmids [31]. As a result, the plasmid carriers combined with the non-carriers, in this study, showed more antibiotic resistance. According to this previous study by Svara et al. these observations could have been a result of horizontal gene transfer where wild-type cells were more normally treated as the plasmids of wild-type bacteria that allow additional plasmids to enter bacterial cells [31]. Therefore,

in a continuous presence of antibiotics, resistance genes in the host genome develop a high tolerance of treatments even in the presence of antibiotics.

Antibiotic resistance presents a pertinent problem for public health [31]. When resistance genes are transported through plasmid carriage or by phages, the spread can accelerate. Plasmids can allow the rapid spread of antibiotic resistance genes [31]. However, many pressures that favor the plasmid carriage of resistance genes have not been fully examined [31]. For plasmid maintenance in a bacterial population, bacteria carrying plasmids and the wild-type cells require the presence of plasmid-free bacterial cells [31]. A reason for this is that plasmid carriage combined with resistance has a cost of fitness that must be balanced with the availability of infection susceptible plasmid-free cells [31]. When the plasmid-free cells are eliminated by antibiotics, the plasmids have fewer cells to infect [31]. Hence, in this study, there may have been a higher fitness cost of the plasmid carriage in the B.sub-C-VH-*E. coli* samples because antibiotic resistance increased. The increased numbers of resistant bacterial cells outcompeted the reduced numbers of plasmid-free bacterial cells, including *V. harveyi* and *E. coli*, and this caused a high cost of fitness. However, Svava et al. showed that plasmids with lower transfer rates cannot carry antibiotic resistance genes [31].

As a consequence, our understanding of the conditions where antibiotic resistant genes are carried in plasmids can assist with developing strategies to lessen the proliferation and spread of resistance via blocking plasmid transfer [31]. Without the presence of antibiotic-mediated selection, bacteria can remove their resistant bacterial competitors when the resistance genes cause a high fitness cost [32-37]. Conjugation and horizontal gene transfer allow for the ample persistence of plasmids, but an extremely high rate of conjugation is needed [38-44]. Conjugative plasmids can be transferred when there is a higher rate of conjugation remaining after the absence of antibiotics in *E. coli* samples [45-51]. The results from the Lopatkin et al. study showed that lessening the use of antibiotics is not sufficient enough for eliminating resistance [32-37]. As a consequence, joining conjugation inhibition with increasing plasmid loss could become an improved method to lessen conjugation-dependent persistence of antibiotic resistance genes [32-37].

4.4. HYPOTHETICAL MODEL FOR AN INTEGRATIVE CYCLE OF PLASMID PERSISTENCE

Plasmids can be shared and carry genes that assist bacterial hosts to survive in adverse and complex environments [1]. Additionally, when plasmids exhibit epistasis (fitness) and interact, the plasmids can be maintained in homeostasis in the population where this combination offers a benefit of fitness to the bacterial host cells [12-17]. Plasmid-free cells can be non-present in the last and final bacterial population [12-17]. Bacterial populations with one favourable plasmid did not have plasmid-free cells lacking in the final population [12-17]. However, advantageous bacteria can still become extinct because only garnering fitness does not guarantee plasmid persistence [12]. Gama et al. results show that different assortments having positive epistasis can only survive and continue to proliferate when there is a lowered fitness cost, elevated rates of conjugation, and a decreased loss of transfer rates [12]. Contrarily, having low fitness cost, high rates of conjugation and reduced rates of loss is not a realistic condition for maintaining plasmid persistence [12-17]. These three conditions are not probable because the rate of conjugation is dependent on the costs of fitness, as a result of increased expenditure of energy as well as metabolites where only plasmids can benefit from conjugation activity at a lower cost, but plasmids cannot persist in both of these two conditions, simultaneously. Thus, according to Roder et al., bacterial cells that carry plasmids increase the removal of bacterial cells lacking plasmids where biofilms can become temporary storages for plasmids [30]. Hence, Figure 6 shows a potential model for a cycle of plasmid persistence. The model is integrative in that it combines the three natural bacterial conditions, examined in this study, with host cell fitness and growth, conjugation, horizontal gene transfer (HGT), and the carriage of plasmids. The model may offer a potential visual explanation for the maintenance of plasmids in natural complex systems such as in the biofilm formed in an abiotic gut microbiome or in a biotic ecosystem.




Figure 6: An integrative and Hypothetical Model for the Cycle of Plasmid Persistence. This hypothetical model shows a cycle for plasmid persistence in natural bacterial populations within communities of biofilm formed on abiotic or biotic surfaces. These bacterial communities begin growing with host cell plasmid carriers and non-carriers. The plasmid carriers grow more vigorously than the non-carriers until there are no plasmid-free bacteria available. At this point, there is a high fitness cost with low conjugation and low horizontal gene transfer (HGT). The plasmid carriers then begin losing plasmids at a high rate, and then the plasmids are stored in the biofilm. This generates more plasmid-free cells where the plasmids can then enter the cells. The level of conjugation heightens with a low fitness cost and a higher rate of HGT. Bacterial cells re-enter a state of growth to restart the cycle of plasmid persistence.

5. CONCLUSIONS

In this study, the natural phases of bacterial growth, biofilm production, and antibiotic resistance were examined in relation to plasmid carriage. These intrinsic stages combined with plasmid carriage were surveyed through *in vitro* co-culture assays in order to form a potential explanation for plasmid persistence in natural conditions. Major changes in these natural states were observed in the samples with the plasmid carriers versus the non-carriers. Bacterial growth decreased in all carrier co-cultures, biofilm was produced in *B. subtilis* carriers with *V. harveyi*, biofilm was reduced in *B. subtilis* carriers with *E. coli*, and the *B. subtilis* carrier *V. harveyi* and *E. coli* co-cultures increased in antibiotic resistance. It was concluded that plasmid carriage affects bacterial growth, biofilm formation, and the rate of antibiotic resistance, in which plasmid carriage, including each altered bacterial phase, contributes to plasmid persistence. However, a major limitation of this present study includes that the effects of plasmid fitness were only examined through *in vitro* methods of LB liquid culturing. This is a common practice in the field where past studies have displayed the effects of plasmid fitness with laboratory conditions that are associated with the same data measured in animal

models [1-5]. As a result, the results in this present study may not completely represent the plasmid fitness in the human gut. Future studies are needed to further survey the total complexity of *in vitro* matrices with focused analysis of *in vivo* animal models. However, analyzing the cost of plasmid production in liquid cultures cannot represent the totality of bacterial dynamics in a spatial ecosystem, therefore, an image of observation created through a mathematical model of plasmid dynamics in bacterial biofilms and colonies is necessary for predicting the outcome of plasmids in bacterial populations [18]. Developing this model can be improved by including three dimensions such as multiple strains and multiple species of bacteria from many different ecosystems [18]. Including more dimensions can enhance our understanding and predictions for plasmids in spatial bacterial communities [18]. Through exploring these multi-dimensional microbial communities, more methods to stagnate the spread of plasmids can be found that may limit the propagation of antibiotic resistance.

6. CONFLICT OF INTERESTS

The authors declare no conflict of interest.

7. ACKNOWLEDGEMENTS

Many thanks are given to my mentors who provided lectures, workshops, seminars, and research readings, which helped to establish my understanding of the fundamentals for microbiology. I also received hands-on experience in applying common and current novel microbiological techniques.

8. REFERENCES

1. Alonso-Del Valle A, León-Sampedro R, Rodríguez-Beltrán J, DelaFuente J, Hernández-García M, Ruiz-Garbajosa P, et al. Variability of plasmid fitness effects contributes to plasmid persistence in bacterial communities. *Nat Commun.* 2021;12(1):2653. doi: 10.1038/s41467-021-22849-y.
2. Pinilla-Redondo R, Olesen AK, Russel J, de Vries LE, Christensen LD, Musovic S, et al. Broad Dissemination of Plasmids across Groundwater-Fed Rapid Sand Filter Microbiomes. *mBio.* 2021;12(6):e0306821. doi: 10.1128/mBio.03068-21.
3. Hall JPJ, Wright RCT, Harrison E, Muddiman KJ, Wood AJ, Paterson S, et al. Plasmid fitness costs are caused by specific genetic conflicts enabling resolution by compensatory mutation. *PLoS Biol.* 2021;19(10):e3001225. doi: 10.1371/journal.pbio.3001225.
4. Jordt H, Stalder T, Kosterlitz O, Ponciano JM, Top EM, Kerr B. Coevolution of host-plasmid pairs facilitates the emergence of novel multidrug resistance. *Nat Ecol Evol.* 2020;4(6):863-9. doi: 10.1038/s41559-020-1170-1.
5. Li L, Dechesne A, Madsen JS, Nesme J, Sørensen SJ, Smets BF. Plasmids persist in a microbial community by providing fitness benefit to multiple phylotypes. *ISME J.* 2020;14(5):1170-81. doi: 10.1038/s41396-020-0596-4.
6. Stecher B, Denzler R, Maier L, Bernet F, Sanders MJ, Pickard DJ, et al. Gut inflammation can boost horizontal gene transfer between pathogenic and commensal Enterobacteriaceae. *Proc Natl Acad Sci U S A.* 2012 Jan 24;109(4):1269-74. doi: 10.1073/pnas.1113246109.
7. Meijer WJJ, Boer DR, Ares S, Alfonso C, Rojo F, Luque-Ortega JR, et al. Multiple Layered Control of the Conjugation Process of the *Bacillus subtilis* Plasmid pLS20. *Front Mol Biosci.* 2021;8:648468. doi: 10.3389/fmolb.2021.648468.
8. Liu L, Chen X, Skogerbø G, Zhang P, Chen R, He S, Huang DW. The human microbiome: a hot spot of microbial horizontal gene transfer. *Genomics.* 2012;100(5):265-70. doi: 10.1016/j.ygeno.2012.07.012.
9. Shterzer N, Mizrahi I. The animal gut as a melting pot for horizontal gene transfer. *Can J Microbiol.* 2015;61(9):603-5. doi: 10.1139/cjm-2015-0049.
10. Sarker MM, Islam KN, Huri HZ, Rahman M, Imam H, Hosen MB, et al. Studies of the impact of occupational exposure of pharmaceutical workers on the development of antimicrobial drug resistance. *J Occup Health.* 2014;56(4):260-70. doi: 10.1539/joh.14-0012-0a.
11. Stalder T, Cornwell B, Lacroix J, Kohler B, Dixon S, Yano H, et al. Evolving Populations in Biofilms Contain More Persistent Plasmids. *Mol Biol Evol.* 2020;37(6):1563-76. doi: 10.1093/molbev/msaa024.
12. Gama JA, Zilhão R, Dionisio F. Plasmid Interactions Can Improve Plasmid Persistence in Bacterial Populations. *Front Microbiol.* 2020;11:2033. doi: 10.3389/fmicb.2020.02033.
13. Hall JPJ, Wright RCT, Guymier D, Harrison E, Brockhurst MA. Extremely fast amelioration of plasmid fitness costs by multiple functionally diverse pathways. *Microbiology (Reading).* 2020;166(1):56-62. doi: 10.1099/mic.0.000862.
14. Hülter NF, Wein T, Effe J, Garoña A, Dagan T. Intracellular Competitions Reveal Determinants of Plasmid Evolutionary Success. *Front Microbiol.* 2020;11:2062. doi: 10.3389/fmicb.2020.02062.
15. Dionisio F, Zilhão R, Gama JA. Interactions between plasmids and other mobile genetic elements affect their transmission and persistence. *Plasmid.* 2019;102:29-36. doi: 10.1016/j.plasmid.2019.01.003.
16. Gama JA, Fredheim EGA, Cléon F, Reis AM, Zilhão R, Dionisio F. Dominance Between Plasmids Determines the Extent of Biofilm Formation. *Front Microbiol.* 2020;11:2070. doi: 10.3389/fmicb.2020.02070.
17. Hernández-Beltrán JCR, San Millán A, Fuentes-Hernández A, Peña-Miller R. Mathematical Models of Plasmid Population Dynamics. *Front Microbiol.* 2021;12:606396. doi: 10.3389/fmicb.2021.606396.
18. Krone SM, Lu R, Fox R, Suzuki H, Top EM. Modelling the spatial dynamics of plasmid transfer and persistence. *Microbiology (Reading).* 2007;153(Pt 8):2803-16. doi: 10.1099/mic.0.2006/004531-0.
19. Benz F, Huisman JS, Bakkeren E, Herter JA, Stadler T, Ackermann M, et al. Plasmid- and strain-specific factors drive variation in ESBL-plasmid spread in vitro and in vivo. *ISME J.* 2021;15(3):862-78. doi: 10.1038/s41396-020-00819-4.
20. Gama JA, Fredheim EGA, Cléon F, Reis AM, Zilhão R, Dionisio F. Dominance Between Plasmids Determines the Extent of Biofilm Formation. *Front Microbiol.* 2020;11:2070. doi: 10.3389/fmicb.2020.02070.
21. Nesse LL, Mo SS, Ramstad SN, Witsø IL, Sekse C, Bruvoll AEE, et al. The Effect of Antimicrobial Resistance Plasmids Carrying bla_{CMY-2} on Biofilm Formation by *Escherichia coli* from the Broiler Production Chain. *Microorganisms.* 2021;9(1):104. doi: 10.3390/microorganisms9010104.
22. Yildiz FH, Visick KL. *Vibrio* biofilms: so much the same yet so different. *Trends Microbiol.* 2009;17(3):109-18. doi: 10.1016/j.tim.2008.12.004.
23. Guttenplan SB, Kearns DB. Regulation of flagellar motility during biofilm formation. *FEMS Microbiol Rev.* 2013;37(6):849-71. doi: 10.1111/1574-6976.12018.
24. Belas R. Biofilms, flagella, and mechanosensing of surfaces by bacteria. *Trends Microbiol.* 2014;22(9):517-27. doi: 10.1016/j.tim.2014.05.002.
25. Wood TK, González Barrios AF, Herzberg M, Lee J. Motility influences biofilm architecture in *Escherichia coli*. *Appl Microbiol Biotechnol.* 2006;72(2):361-7. doi: 10.1007/s00253-005-0263-8.
26. Pesavento C, Becker G, Sommerfeldt N, Possling A, Tschowri N, Mehlis A. Inverse regulatory coordination of motility and curli-mediated adhesion in *Escherichia coli*. *Genes Dev.* 2008;22(17):2434-46. doi: 10.1101/gad.475808.
27. Hölscher T, Bartels B, Lin YC, Gallegos-Monterrosa R, Price-Whelan A, Kolter R, et al. Motility, Chemotaxis and Aerotaxis Contribute to Competitiveness during Bacterial Pellicle Biofilm Development. *J Mol Biol.* 2015;427(23):3695-708. doi: 10.1016/j.jmb.2015.06.014.
28. Haiko J, Westerlund-Wikström B. The role of the bacterial flagellum in adhesion and virulence. *Biology (Basel).* 2013;2(4):1242-67. doi: 10.3390/biology2041242.
29. Connelly MB, Young GM, Sloma A. Extracellular proteolytic activity plays a central role in swarming motility in *Bacillus subtilis*. *J Bacteriol.* 2004;186(13):4159-67. doi: 10.1128/JB.186.13.4159-4167.2004.
30. Røder HL, Trivedi U, Russel J, Kragh KN, Herschend J, Thalsø-Madsen I, et al. Biofilms can act as plasmid reserves in the absence of plasmid specific selection. *NPJ Biofilms Microbiomes.* 2021;7(1):78. doi: 10.1038/s41522-021-00249-w.
31. Svava F, Rankin DJ. The evolution of plasmid-carried antibiotic resistance. *BMC Evol Biol.* 2011;11:130. doi: 10.1186/1471-2148-11-130.
32. Lopatkin AJ, Meredith HR, Srimani JK, Pfeiffer C, Durrett R, You L. Persistence and reversal of plasmid-mediated antibiotic resistance. *Nat Commun.* 2017;8(1):1689. doi: 10.1038/s41467-017-01532-1.
33. San Millan A. Evolution of Plasmid-Mediated Antibiotic Resistance in the Clinical Context. *Trends Microbiol.* 2018;26(12):978-85. doi: 10.1016/j.tim.2018.06.007.
34. San Millan A, Santos-Lopez A, Ortega-Huedo R, Bernabe-Balas C, Kennedy SP, Gonzalez-Zorn B. Small-plasmid-mediated antibiotic resistance is enhanced by increases in plasmid copy number and bacterial fitness. *Antimicrob Agents Chemother.* 2015;59(6):3335-41. doi: 10.1128/AAC.00235-15.
35. Andersson DI, Hughes D. Persistence of antibiotic resistance in bacterial populations. *FEMS Microbiol Rev.* 2011;35(5):901-11. doi: 10.1111/j.1574-6976.2011.00289.x.
36. Loftie-Eaton W, Bashford K, Quinn H, Dong K, Millstein J, Hunter S, et al. Compensatory mutations improve general permissiveness to antibiotic resistance plasmids. *Nat Ecol Evol.* 2017;1(9):1354-63. doi: 10.1038/s41559-017-0243-2.
37. Harrison E, Guymier D, Spiers AJ, Paterson S, Brockhurst MA. Parallel compensatory evolution stabilizes plasmids across the parasitism-mutualism continuum. *Curr Biol.* 2015;25(15):2034-9. doi: 10.1016/j.cub.2015.06.024.
38. Kottara A, Hall JPJ, Harrison E, Brockhurst MA. Variable plasmid fitness effects and mobile genetic element dynamics across *Pseudomonas* species. *FEMS Microbiol Ecol.* 2018;94(1):fix172. doi: 10.1093/femsec/fix172.
39. Bottery MJ, Wood AJ, Brockhurst MA. Adaptive modulation of antibiotic resistance through intragenomic coevolution. *Nat Ecol Evol.* 2017;1(9):1364-69. doi: 10.1038/s41559-017-0242-3.
40. Ilhan J, Kupczok A, Woehle C, Wein T, Hülter NF, Rosenstiel P, et al. Segregational Drift and the Interplay between Plasmid Copy Number and Evolvability. *Mol Biol Evol.* 2019;36(3):472-86. doi: 10.1093/molbev/msy225.

41. Harrison E, Brockhurst MA. Plasmid-mediated horizontal gene transfer is a coevolutionary process. *Trends Microbiol.* 2012;20(6):262-7. doi: 10.1016/j.tim.2012.04.003.
42. San Millan A, Escudero JA, Gifford DR, Mazel D, MacLean RC. Multicopy plasmids potentiate the evolution of antibiotic resistance in bacteria. *Nat Ecol Evol.* 2016;1(1):10. doi: 10.1038/s41559-016-0010.
43. Turner PE, Williams ES, Okeke C, Cooper VS, Duffy S, Wertz JE. Antibiotic resistance correlates with transmission in plasmid evolution. *Evolution.* 2014;68(12):3368-80. doi: 10.1111/evo.12537.
44. Santos-Lopez A, Bernabe-Balas C, Ares-Arroyo M, Ortega-Huedo R, Hoefler A, San Millan A, et al. A Naturally Occurring Single Nucleotide Polymorphism in a Multicopy Plasmid Produces a Reversible Increase in Antibiotic Resistance. *Antimicrob Agents Chemother.* 2017;61(2):e01735-16. doi: 10.1128/AAC.01735-16.
45. Wein T, Hülter NF, Mizrahi I, Dagan T. Emergence of plasmid stability under non-selective conditions maintains antibiotic resistance. *Nat Commun.* 2019;10(1):2595. doi: 10.1038/s41467-019-10600-7.
46. Porse A, Schønning K, Munck C, Sommer MO. Survival and Evolution of a Large Multidrug Resistance Plasmid in New Clinical Bacterial Hosts. *Mol Biol Evol.* 2016;33(11):2860-73. doi: 10.1093/molbev/msw163.
47. San Millan A, Peña-Miller R, Toll-Riera M, Halbert ZV, McLean AR, Cooper BS, et al. Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids. *Nat Commun.* 2014;5:5208. doi: 10.1038/ncomms6208.
48. Zwanzig M, Harrison E, Brockhurst MA, Hall JPJ, Berendonk TU, Berger U. Mobile Compensatory Mutations Promote Plasmid Survival. *mSystems.* 2019;4(1):e00186-18. doi: 10.1128/mSystems.00186-18.
49. Rodriguez-Beltran J, Hernandez-Beltran JCR, DelaFuente J, Escudero JA, Fuentes-Hernandez A, MacLean RC, et al. Multicopy plasmids allow bacteria to escape from fitness trade-offs during evolutionary innovation. *Nat Ecol Evol.* 2018;2(5):873-881. doi: 10.1038/s41559-018-0529-z.
50. San Millan A, Toll-Riera M, Qi Q, Betts A, Hopkinson RJ, McCullagh J, et al. Integrative analysis of fitness and metabolic effects of plasmids in *Pseudomonas aeruginosa* PAO1. *ISME J.* 2018;12(12):3014-24. doi: 10.1038/s41396-018-0224-8.
51. San Millan A, MacLean RC. Fitness Costs of Plasmids: a Limit to Plasmid Transmission. *Microbiol Spectr.* 2017;5(5). doi: 10.1128/microbiolspec.MTBP-0016-2017.