


Original article

Healthcare Providers' Compliance with Guidelines for Catheter-Associated Urinary Tract Infections in a Rural Teaching and Referral Hospital

Reha Girgin ^{a,*} , Emre Horuz ^b

^a Zonguldak Bulent Ecevit University Hospital, Department of Urology, Turkey

^b Zonguldak Bulent Ecevit University Hospital, Department of Infectious Diseases, Turkey

ARTICLE INFO
Article history:

Received 06 July 2022

Received in revised form 23

August 2022

Accepted 02 September 2022

Keywords:

Urinary tract infection

Catheter

Health personnel

Knowledge

Prevention

ABSTRACT

Introduction: Healthcare professionals (HCPs) play a key role in the prevention of catheter-associated urinary tract infections (CAUTI). This study aims to determine the compliance of healthcare personnel with urethral catheterisation standards and the factors affecting it.

Materials and Methods: This study is a prospective survey and was conducted between February and March 2021 after ethical approval and patient consent. An internet-based, structured 39-item questionnaire was sent via email to all HCPs at Zonguldak Bulent Ecevit University Hospital, Zonguldak, Turkey. A to evaluate their knowledge and attitudes towards the prevention of CAUTI. The questionnaire was completed by 156 health professionals. For categorical parameters Fisher's exact test, though for scale parameters student t-test, Mann-Whitney U and Kruskal-Wallis tests were used. All statistical analysis was done using SPSS 17.0.

Results: The average years of experience of the HCPs who participated in the survey was 12.5 years. The level of knowledge of the doctors and the nurses about the indication of catheter insertion was similar, whereas the level of knowledge about CAUTI prevention was higher among the doctors. However, none of the participants could fully describe how to prevent CAUTI.

Conclusions: The level of knowledge of CAUTI preventive measures among the participants was insufficient. To effectively preventing CAUTI, there is a need to change HCPs' perspective on this issue, which can be achieved through training, and advantage current technologies.

© 2022 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author.

E-mail address: mujdereha7477@hotmail.com

ISSN: 2695-5075 / © 2022 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.53986/ibjm.2022.0036>

Cumplimiento de las pautas para las infecciones del tracto urinario asociadas con el catéter por parte de los proveedores de atención médica en un hospital rural docente y de referencia

INFO. ARTÍCULO

Historia del artículo:

Recibido 06 Julio 2022

Recibido en forma revisada 23

Agosto 2022

Aceptado 02 Septiembre 2022

Palabras clave:

Infecciones del tracto urinario

Catéter

Personal sanitario

Conocimiento

Prevención

RESUMEN

Introducción: Los profesionales de la salud (PS) juegan un papel clave en la prevención de infecciones del tracto urinario asociadas al catéter (ITUAC). Este estudio tiene como objetivo determinar el cumplimiento de las normas de sondaje uretral por parte del personal sanitario y los factores que lo afectan.

Materiales y métodos: Este estudio es una encuesta prospectiva y se realizó entre febrero y marzo de 2021 después de la aprobación ética y el consentimiento del paciente. Se envió un cuestionario estructurado de 39 ítems basado en Internet por correo electrónico a todos los HCP en el Hospital Universitario Zonguldak Bulent Ecevit, Zonguldak, Turquía. A para evaluar sus conocimientos y actitudes hacia la prevención de la ITUAC. El cuestionario fue completado por 156 profesionales de la salud. Para los parámetros categóricos se utilizó la prueba exacta de Fisher, aunque para los parámetros de escala se utilizaron la prueba t de Student, la U de Mann-Whitney y la de Kruskal-Wallis. Todos los análisis estadísticos se realizaron con SPSS 17.0.

Resultados: El promedio de años de experiencia de los profesionales de la salud que participaron en la encuesta fue de 12,5 años. El nivel de conocimiento de los médicos y las enfermeras sobre la indicación de la inserción del catéter fue similar, mientras que el nivel de conocimiento sobre la prevención de las ITUAC fue mayor entre los médicos. Sin embargo, ninguno de los participantes pudo describir completamente cómo prevenir ITUAC.

Conclusiones: El nivel de conocimiento de las medidas preventivas de ITUAC entre los participantes fue insuficiente. Para prevenir CAUTI de manera efectiva, es necesario cambiar la perspectiva de los profesionales de la salud sobre este tema, lo que se puede lograr a través de la capacitación y aprovechar las tecnologías actuales.

© 2022 Los Autores. Publicado por Iberoamerican Journal of Medicine. Éste es un artículo en acceso abierto bajo licencia CC BY (<http://creativecommons.org/licenses/by/4.0/>).

HOW TO CITE THIS ARTICLE: Girgin R, Horuz E. Healthcare Providers' Compliance with Guidelines for Catheter-Associated Urinary Tract Infections in a Rural Teaching and Referral Hospital. *Iberoam J Med.* 2022;4(4):199-205. doi: 10.53986/ibjm.2022.0036.

1. INTRODUCTION

Nosocomial infections are very common, and approximately 40% of these infections are catheter-related urinary tract infections (CAUTI) [1]. A Foley catheter, which is the most important predisposing factor of CAUTI, is used in 25% of hospitalised patients generally and almost 100% of patients in intensive care [1].

CAUTI is an important cause of morbidity and mortality in hospitalised patients [2]. It also creates a serious reason for the irrational use of antibiotics and the development of resistant microorganisms, putting other patients at risk of being infected with these resistant organisms [1-3]. CAUTI increases susceptibility to catheter occlusion in catheterised patients and predisposition to urinary system stones, including malignant transformation [1]. This susceptibility, paradoxically, also increases the risk of CAUTI [1].

The risk of CAUTI increases with each passing day with a

catheter [1]. The incidence of CAUTI has been found to be similar in patients receiving intensive and nonintensive care [4]. The most important parameter to prevent CAUTI is to avoid unnecessary catheterisation; thus, it is important to evaluate all patients with a catheter [5]. Although infection control committees in hospitals have established strict follow-up protocols regarding CAUTI, if doctors and nurses have sufficient knowledge of this issue, it will help in reducing such infections [6]. In this study, we aimed to determine the compliance of healthcare personnel with urethral catheterisation standards and the factors affecting it in a rural academic hospital in Turkey.

2. MATERIAL AND METHODS

Ethical approval for this study was obtained from Zonguldak Bulent Ecevit University Local Ethics Committee (meeting date: 10/02/2021; meeting no: 2021/03). A questionnaire

was emailed to 712 healthcare professionals (HCPs) (330 doctors at different academic levels and 382 nurses) and asked to respond after obtaining their written consent. The present research study included participants working in different departments and areas (i.e. intensive care, operating room, ward etc.) in Zonguldak Bulent Ecevit University Hospital, Zonguldak, Turkey between February and March 2021. Google Docs™ was used to prepare the survey form, as it ensured the participants' anonymity. Furthermore, the participants were informed that the results of the questionnaire would be used for scientific purposes only.

before performing the analysis agreeing the method employed by Jain et al [1]. The values were expressed as follows: (4): Almost always indicated; (3): Sometimes indicated as indicated; (2): Rarely indicated; (1): Never indicated as not indicated) (4: Large effect; 3: Moderate effect as effective; 2: Some effect; 1: No effect as ineffective).

2.1. STATISTICAL METHODS

Nominal and ordinal parameters were reported as frequencies. Fischer's exact test was used to assess the

Table 1: Baseline characteristics of study group and difference analysis results

	Doctors (n=82) n (%)	Nurses (n=74) n (%)	Total (n=156) n (%)	P value
Age	39.84±9.54	34.57±6.06	37.34±8.48	0.000
Gender				
Males	51 (62.2)	24 (32.4)	75 (48.1)	0.000
Females	31 (37.8)	50 (67.6)	81 (51.9)	
Department				
Service/Polyclinic	41 (50.0)	52 (70.3)	93 (59.6)	0.000
Intensive Care	5 (6.1)	18 (24.3)	23 (14.7)	
Operating Room	36 (43.9)	4 (5.4)	40 (25.6)	
Branch				
Internal	30 (36.6)	44 (59.5)	74 (47.4)	0.003
Surgery	52 (63.4)	30 (40.5)	82 (52.6)	
Experience	12.73±9.38	12.27±6.06	12.51±7.96	0.844
Academic position				
Professor	19 (23.2)	-	19 (12.2)	0.000
Associate Professor	14 (17.1)	-	14 (9.0)	
Assistant Professor	20 (24.4)	-	20 (12.8)	
Senior Assistant	18 (21.9)	-	18 (11.5)	
Assistant under 2 years in experience	11 (13.4)	-	11 (7.1)	
Nurse In Charge	-	20 (27.1)	20 (12.8)	
Clinic nurse	-	54 (72.9)	54 (34.6)	

The survey questions for this present study were adapted particularly for this study from the questions prepared according to the Centers for Disease Control guidelines by Jain et al [1, 7]. The authors to ensure they reflect the knowledge and skills of the target audience and that they are simple and understandable previously evaluated the questions. Demographic data (such as age, gender, academic position, department and area of posting) of the participants were collected during the survey. The second section of the survey had 10 questions that evaluated the participants' knowledge of the indication for catheter insertion, the third part had 15 questions that assessed their knowledge level about CAUTI prevention, and the last part had 8 questions that evaluated the general attitudes of the participants /towards CAUTI.

The participants were asked to answer questions on the indication for catheter insertion and methods of preventing CAUTI on a four-point scale. After comparing the answers with the Centers for Disease Control guidelines, the questionnaire responses were converted to binary variables

differences between the categorical parameters. Means and standard deviations were used for the description of the scale parameters. The Kolmogorov–Smirnov test was used to evaluate the normality of the scale parameters. The independent samples t-test was used for the normally distributed parameters, while the Mann–Whitney U and Kruskal–Wallis tests were used for no normally distributed parameters. All the analyses were performed using SPSS 17.0 for Windows, with a 95% confidence interval and 0.05 significance level.

3. RESULTS

Of the 712 HCPs in the hospital, 156 (21.91%) participated in the survey. The mean age of the participants was 37.34 ± 8.48, and 48.1% of the participants were male. Of the 156 HCPs, 82 (52.6%) were doctors, while 74 (47.4%) were nurses. The baseline characteristics of the participants are summarised in Table 1.

Table 2: Frequency of respondents having identified the indications correctly

	Doctors (n=82) n (%)	Nurses (n=74) n (%)	Total (n=156) n (%)	P value
Urethral stricture causing obstruction to urinary flow	65 (79.3)	53 (71.6)	118 (75.6)	0.178
Neurogenic bladder due to paraplegia or quadriplegia	68 (82.9)	54 (73.0)	122 (78.2)	0.095
Prolonged immobilization due to unstable lumbar spine fracture	60 (73.2)	55 (74.3)	115 (73.7)	0.508
Urine output monitoring in a mobile patient	59 (72.0)	45 (60.8)	104 (66.7)	0.096
For assisting healing of decubitus ulcers in incontinent patients	56 (68.3)	56 (75.7)	112 (71.8)	0.199
Obtaining urine sample for culture and sensitivity testing	48 (58.5)	41 (55.4)	89 (57.1)	0.408
Palliative care in terminally ill patient	67 (81.7)	60 (81.1)	127 (81.4)	0.541
Nursing care for incontinent patient	37 (45.1)	22 (29.7)	59 (37.8)	0.034
Routinely before any kind of surgical procedure in a patient	26 (31.7)	22 (29.7)	48 (30.8)	0.463
In patients anticipated to receive large volume infusions or diuretics during surgery	65 (79.3)	59 (79.7)	124 (79.5)	0.551

Regarding the participants' level of knowledge of the indications for urethral catheterisation, there was a significant difference between doctors and nurses' knowledge of nursing care for incontinent patients ($p < 0.05$). All answers to the other questions were statistically significant ($p > 0.05$) (Table 2).

infected patients', 'Prophylactic antimicrobials should be given for three days when a catheter is inserted' and 'Routinely using antimicrobial coated catheters' were statistically significant between the doctors and the nurses ($p < 0.05$). All other answers given to questions regarding indications for preventing CAUTI were not statistically

Table 3: Number of doctors and nurses correctly identifying methods to prevent CAUTI

	Doctors (n=82) n (%)	Nurses (n=74) n (%)	Total (n=156) n (%)	P value
Hand hygiene should be done immediately before and after any manipulation of catheter site or apparatus	77 (93.9)	72 (97.3)	149 (95.5)	0.266 ^a
As small a catheter as possible should be used to minimize urethral trauma	59 (72.0)	54 (73.0)	113(72.4)	0.515 ^a
It should be inserted only when necessary and removed as soon as possible	82 (100.0)	73 (98.6)	155 (99.4)	0.474 ^a
Use of other methods of urinary drainage such as condom catheter drainage, suprapubic or intermittent catheterization for selected patients	3 (3.7)	2 (2.7)	5 (3.2)	0.549 ^a
Avoid kinking of the catheter to maintain an unobstructed flow of urine	78 (95.1)	71 (95.9)	149 (95.5)	0.557 ^a
Irrigation of the bladder with antimicrobial solution/iodine solution at least once daily	59 (72.0)	35 (47.3)	94 (60.3)	0.001 ^a
Twice daily meatal care with antiseptic solution	27 (32.9)	13 (17.6)	40 (25.6)	0.022 ^a
Collecting bag should be emptied regularly	75 (91.5)	67 (90.5)	142 (91.0)	0.530 ^a
Collecting bag should be kept below the level of the bladder	76 (92.7)	72 (97.3)	148 (94.9)	0.174 ^a
Regular bacteriological monitoring of catheterized patients	33 (40.2)	10 (13.5)	43 (27.6)	0.000 ^a
Catheter should be inserted only by personnel proficient in technique of aseptic insertion	72 (87.8)	72 (97.3)	144 (92.3)	0.025 ^a
Isolation of patients known to have UTI from other no infected patients	48 (58.5)	30 (40.5)	78 (50.0)	0.018 ^a
Prophylactic antimicrobials should be given for 3 days when catheter is inserted	65 (79.3)	37 (50.0)	102 (65.4)	0.000 ^a
Regular educational training regarding basic urinary catheter care	78 (95.1)	70 (94.6)	148 (94.9)	0.582 ^a
Routinely using antimicrobial coated catheters	38 (46.3)	46 (62.2)	84 (53.8)	0.034 ^a
Hand hygiene should be done immediately before and after any manipulation of catheter site or apparatus	77 (93.9)	72 (97.3)	149 (95.5)	0.266 ^a
As small a catheter as possible should be used to minimize urethral trauma	59 (72.0)	54 (73.0)	113(72.4)	0.515 ^a
It should be inserted only when necessary and removed as soon as possible	82 (100.0)	73 (98.6)	155 (99.4)	0.474 ^a

The answers given to 'Irrigation of the bladder with an antimicrobial solution/iodine solution at least once daily', 'Twice daily metal care with an antiseptic solution', 'Regular bacteriological monitoring of catheterised patients', 'Catheter should be inserted only by personnel proficient in the technique of aseptic insertion', 'Isolation of patients known to have urinary tract infection from other no

different ($p > 0.05$) (Table 3).

The knowledge level on indications for catheterisation was not significantly different between occupation, gender, position, department, branch and experiment groups ($p > 0.05$). The knowledge level on prevention of CAUTI was significantly different between occupation and academic positions ($p < 0.05$), with the doctors having higher scores

(Table 4).

Among the questions about the attitudes of the doctors and the nurses towards urinary catheterisation practices, the answer given to the item 'Catheter can be inserted for nursing staff convenience' was statistically significant ($p < 0.05$). No significant difference was found between all other responses ($p > 0.05$) (Table 5).

compliance with these rules can reduce the rate of CAUTI by almost one-third [5]. CAUTI rates are thought to be higher than actually reported [8]. Although many factors may play a role in the development of CAUTI, it is known that the most important is the duration of catheterisation. The daily risk for infective consequences is thought to be 5%–10%, which approaches 100% at the end of a monthly

Table 4: Univariate predictors of indications of catheterization

	Indication		Prevention	
	Mean±SD	P value	Mean±SD	P value
Occupation				
Doctor	6.72±1.42	0.404	10.61±1.57	0.000
Nurses	6.31±1.98		9.78±1.34	
Gender				
Males	6.41±1.64	0.245	10.21±1.70	0.839
Females	6.63±1.78		10.22±1.33	
Position				
Lecturer	6.44±1.44	0.067	10.61±1.48	0.001
Assistant doctor	7.25±1.24		10.61±1.75	
Nurse	6.31±1.98		9.78±1.34	
Academic position				
Professor	6.32±1.29	0.169	10.63±1.42	0.006
Associate Professor	5.86±1.75		11.07±1.21	
Assistant Professor	7.10±1.25		10.25±1.25	
Senior Assistant	7.17±1.89		9.67±2.00	
Assistant under 2 years in experience	6.73±1.49		10.91±1.38	
Nurse In Charge	6.25±1.74		10.65±1.18	
Clinic nurse	6.41±1.90	9.72±1.48		
Department				
Service/Polyclinic	6.49±1.83	0.262	10.22±1.56	0.778
Intensive Care	6.17±1.61		10.09±1.20	
Operating Room	6.80±1.45		10.27±1.60	
Branch				
Internal	6.43±1.81	0.547	10.28±1.59	0.526
Surgery	6.61±1.63		10.16±1.45	
Experience				
<20 years	6.56±1.79	0.451	10.20±1.46	0.569
≥20 years	6.41±1.42		10.29±1.73	

4. DISCUSSION

To prevent CAUTI, the entire process from insertion to withdrawal should be followed within the framework of the guidelines. It has been shown in the literature that

period [8]. However, as supported by the literature, the use of urinary catheters in hospitalised patients continues even after the indication for use has expired [1, 9]. To prevent CAUTI, which has an important place among hospital-acquired infections, it is necessary to reduce the duration of catheterisation, and HCPs should have sufficient knowledge of this [10].

Table 5: Attitudes of doctors and nurses regarding urinary catheterization practices

	Attitude	Doctors (n=82) n (%)	Nurses (n=74) n (%)	Total (n=156) n (%)	P value
Renewal reminders for catheters prevents CAUTI	Agree	78 (95.1)	67 (90.5)	145 (92.9)	0.211 ^a
Catheter can be inserted for nursing staff convenience	Disagree	80 (97.6)	66 (89.2)	146 (93.6)	0.034 ^a
It helps if CAUTI prevention is in high priority list of hospitals	Agree	59 (72.0)	55 (74.3)	114 (73.1)	0.440 ^a
CAUTI not a very serious illness	Disagree	76 (92.7)	67 (90.5)	143 (91.7)	0.422 ^a
Education regarding basic catheter care helps prevent CAUTI	Agree	78 (95.1)	67 (90.5)	145 (92.9)	0.211 ^a
Catheter should be removed whenever it is convenient for HCP	Disagree	6 (7.3)	7 (9.5)	13 (8.3)	0.422 ^a
CAUTI is a common problem and virtually impossible to prevent it	Disagree	77 (93.9)	63 (85.1)	140 (89.7)	0.062 ^a
Maintaining a closed drainage system prevents CAUTI	Agree	60 (73.2)	55 (74.3)	115 (73.7)	0.508 ^a

It is important to determine the correct indications to prevent unnecessary catheter usage. In general, the knowledge of the doctors and the nurses was similar regarding determining the indications for catheterisation ($p = 0.404$). However, when the preventive methods were compared, it was observed that the level of knowledge of the doctors was higher than that of the nurses ($p < 0.05$). Among the nurses, it was found that nurses-in-charge have more knowledge on preventive methods. Considering that clinical nurses have a more active role in patient care, this issue draws attention. Ensuring the knowledge level of nurses is high will help in determining unnecessary catheterisation indications effectively and promote catheter hygiene while nurses are monitoring patients [1, 11, 12].

Approximately 70.3% of the nurses and 54.6% of the doctors stated that the use of a catheter might be necessary for the nursing care of patients with urinary incontinence. Although the level of knowledge of the doctors on this subject seems to be higher, they responded at a lower rate than expected. When the answers given for other indications were examined, the present study revealed that more than half of the nurses and the doctors gave similar answers.

Only one-third of the nurses and the doctors agreed on the need for routine catheter insertion before any surgery, which is more than expected. Although the current study could not show a significant difference between the knowledge levels of the nurses and the doctors of different degrees about urinary catheterisation indication in the univariate analysis, the knowledge levels of assistant professors and assistant doctors with more than two years of experience seemed to be higher in our study. This may be related to the fact that doctors at this level are constantly reviewing their knowledge. Today, where the urethral catheter is widely used, the correct approach will be to review the catheterisation indications and keep the knowledge levels of doctors and nurses up to date always.

Both physicians and nurses have largely responded favourably to the use of alternative techniques (such as condom catheter drainage, suprapubic or intermittent catheterisation) in suitable patients instead of urethral catheterisation. According to the literature, the fact that catheterisation techniques are seen as alternatives to each other in terms of CAUTI prevention can be considered in terms of preventing some complications, but it does not change the infective results [13]. In this regard, it would be beneficial to provide training that will change the perspectives of healthcare professionals.

Considering the working field and departments, we could not see a relationship among the HCPs regarding catheterisation approaches. However, Tabrisi et al., reported

in their study that HCPs working in surgical wards were better in compliance with catheter care standards [8]. This difference may have been due to the study design because, while our study focused on the level of knowledge, the related study conducted an application-oriented evaluation. One-third of the doctors and almost half of the nurses stated that irrigation of the bladder with antimicrobial solution/iodine solution at least once daily can prevent CAUTI, and 67.4% of doctors and 82.4% of the nurses stated that daily meatal care with an antiseptic solution could prevent the development of CAUTI. Furthermore, 59.8% of the doctors and 86.5% of the nurses reported that taking regular culture samples from catheterised patients is a preventive approach. As can be understood from these results, although not effective, HCPs have developed habits that can both increase health expenditure and unnecessary workload, which is consistent with the literature¹. While simple precautions are sufficient, complex behaviours of HCPs can be overcome with sufficient training.

In the present study, we could not find a relationship between gender differences and the preventive measures of CAUTI. The absence of gender dominance among HCPs in our country seems to explain this situation. Professional experience did not contribute as a preventive factor, which was consistent with the literature [1, 11, 14]. While it was effective for one parameter where the information is variable, it seemed ineffective for another parameter. In light of evidence-based medicine, it is necessary to renew the knowledge of HCPs from different perspectives.

Of the HCPs, 73.1% reported that if hospitals put CAUTI on a high priority list, it could help to prevent CAUTI, and 92.2% reported that using a catheter reminder could be effective in preventing CAUTI. Although there is an effective infection control committee in our hospital, the use of a catheter reminder is not in practice. Similar problems have been reported repeatedly in the literature [1, 15, 16].

A few of the HCPs (6.4%) stated that the urethral catheter can be inserted for the convenience of the nurses, and most of those who gave this answer were nurses. Similar results have been reported in the literature [17], and people who were primarily concerned with the follow-up of catheterised patients gave this response. Such a rate does not seem surprising since nurses are competent in this regard in our hospital as well.

About 8.3% of the HCPs think that CAUTI is not a serious disease, and 10.3% think that CAUTI is an unavoidable problem. This finding is consistent with the literature [1]. In addition, 7.1% of the HCPs think that catheterisation training cannot prevent infection. About 10% of the healthcare workers have serious concerns about catheter-

related infections, and perhaps this reflects their behavioural approaches.

The main limitation of this study is the small number of participants who completed the survey. The low number may be related to the fact that the internet is not a preferred tool. However, this survey further highlighted the lack of compliance of HCPs with catheterisation guidelines. Reaching the participants online was another limitation. However, our approach enabled anonymity and gave participants ample time to reflect on their answers rather than give an immediate response. Another limitation could concern obtaining data directly from individuals in this study. As there may be an incompatibility between application and knowledge, it would be more appropriate to support our finding with an observational study. Since our research was conducted in a single urban, academic medical centre, a multicenter study with large participation will enable us to reach a more reliable conclusion.

5. CONCLUSIONS

The present study finding revealed that the HCPs were incapable of catheter applications. In our practice also, expanding the indications for catheterization in line with patient demands and unnecessarily prolonging the duration of the catheter instead of using alternative urine collection products might be seen. It seems that education on its own is insufficient to correct this challenge. Presently, we think it would be wiser to use computer-based artificial intelligence for tracking catheterisation time, revising of urethral catheterisation indications and producing alternative products for urine collection, considering the annual costs spent for CAUTI.

6. CONFLICT OF INTERESTS

The authors have no conflict of interest to declare. The authors declared that this study has received no financial support.

7. REFERENCES

1. Jain M, Dogra V, Mishra B, Thakur A, Loomba PS. Knowledge and attitude of doctors and nurses regarding indication for catheterization and prevention

of catheter-associated urinary tract infection in a tertiary care hospital. *Indian J Crit Care Med.* 2015;19(2):76-81. doi: 10.4103/0972-5229.151014.

2. Warren JW. The catheter and urinary tract infection. *Med Clin North Am.* 1991;75(2):481-93. doi: 10.1016/s0025-7125(16)30465-5.

3. Getliffe K, Newton T. Catheter-associated urinary tract infection in primary and community health care. *Age Ageing.* 2006;35(5):477-81. doi: 10.1093/ageing/af1052.

4. Lewis SS, Knelson LP, Moehring RW, Chen LF, Sexton DJ, Anderson DJ. Comparison of non-intensive care unit (ICU) versus ICU rates of catheter-associated urinary tract infection in community hospitals. *Infect Control Hosp Epidemiol.* 2013;34(7):744-7. doi: 10.1086/671000.

5. Datta P, Rani H, Chauhan R, Gombar S, Chander J. Health-care-associated infections: Risk factors and epidemiology from an intensive care unit in Northern India. *Indian J Anaesth.* 2014;58(1):30-5. doi: 10.4103/0019-5049.126785.

6. Fakh MG, Dueweke C, Meisner S, Berriel-Cass D, Savoy-Moore R, Brach Net al. Effect of nurse-led multidisciplinary rounds on reducing the unnecessary use of urinary catheterization in hospitalized patients. *Infect Control Hosp Epidemiol.* 2008;29(9):815-9. doi: 10.1086/589584.

7. Hooton TM, Bradley SF, Cardenas DD, Colgan R, Geerlings SE, Rice JC, et al. Diagnosis, prevention, and treatment of catheter-associated urinary tract infection in adults: 2009 International Clinical Practice Guidelines from the Infectious Diseases Society of America. *Clin Infect Dis.* 2010;50(5):625-63. doi: 10.1086/650482.

8. Taleschian-Tabrizi N, Farhadi F, Madani N, Mokhtarkhani M, Kolahdoustan K, Hajebrahimi S. Compliance With Guideline Statements for Urethral Catheterization in an Iranian Teaching Hospital. *Int J Health Policy Manag.* 2015;4(12):805-11. doi: 10.15171/ijhpm.2015.128.

9. Apisarnthanarak A, Rutjanawech S, Wichansavakun S, Ratanabunjerdkul H, Patthranitima P, Thongphubeth K, et al. Initial inappropriate urinary catheters use in a tertiary-care center: incidence, risk factors, and outcomes. *Am J Infect Control.* 2007;35(9):594-9. doi: 10.1016/j.ajic.2006.11.007.

10. Atkins L, Sallis A, Chadborn T, Shaw K, Schneider A, Hopkins S, et al. Reducing catheter-associated urinary tract infections: a systematic review of barriers and facilitators and strategic behavioural analysis of interventions. *Implement Sci.* 2020;15(1):44. doi: 10.1186/s13012-020-01001-2.

11. Apisarnthanarak A, Thongphubeth K, Sirinvaravong S, Kitkangvan D, Yuekyen C, Warachan B, et al. Effectiveness of multifaceted hospitalwide quality improvement programs featuring an intervention to remove unnecessary urinary catheters at a tertiary care center in Thailand. *Infect Control Hosp Epidemiol.* 2007;28(7):791-8. doi: 10.1086/518453.

12. Elpern EH, Killeen K, Ketchem A, Wiley A, Patel G, Lateef O. Reducing use of indwelling urinary catheters and associated urinary tract infections. *Am J Crit Care.* 2009;18(6):535-41; quiz 542. doi: 10.4037/ajcc2009938.

13. Chapple A, Prinjha S, Feneley R. Comparing transurethral and suprapubic catheterization for long-term bladder drainage: a qualitative study of the patients' perspective. *J Wound Ostomy Continence Nurs.* 2015;42(2):170-5. doi: 10.1097/WON.0000000000000096.

14. Drekonja DM, Kuskowski MA, Johnson JR. Internet survey of Foley catheter practices and knowledge among Minnesota nurses. *Am J Infect Control.* 2010;38(1):31-7. doi: 10.1016/j.ajic.2009.05.005.

15. Saint S, Kowalski CP, Forman J, Damschroder L, Hofer TP, Kaufman SR, et al. A multicenter qualitative study on preventing hospital-acquired urinary tract infection in US hospitals. *Infect Control Hosp Epidemiol.* 2008;29(4):333-41. doi: 10.1086/529589.

16. Saint S, Kowalski CP, Kaufman SR, Hofer TP, Kauffman CA, Olmsted RN, et al. Preventing hospital-acquired urinary tract infection in the United States: a national study. *Clin Infect Dis.* 2008;46(2):243-50. doi: 10.1086/524662.

17. Viswanathan K, Rosen T, Mulcare MR, Clark S, Hayes J, Lachs MS, et al. Emergency Department Placement and Management of Indwelling Urinary Catheters in Older Adults: Knowledge, Attitudes, and Practice. *J Emerg Nurs.* 2015;41(5):414-22. doi: 10.1016/j.jen.2015.02.013.