


Editorial

Use of polypill to improve therapeutic adherence and outcomes in adults with recent acute myocardial infarction: is it an effective strategy?

Yelson Alejandro Picón-Jaimes ^{a,*} , María Margarita Garcés-Lince ^b , Juan Fernando Gómez-Gómez ^c , Omar Villamizar-Rueda ^d 

^a Medical and Surgical Research Center, Future Surgeons Chapter, Colombian Surgery Association, Bogotá, Colombia

^b Department of Medicine, Fundación Universitaria San Martín, Pasto, Colombia

^c Department of Medicine, Universidad Remington, Medellín, Colombia

^d Department of Medicine, Universidad de Santander, Bucaramanga, Colombia

ARTICLE INFO
Article history:

Received 24 October 2022

Received in revised form 19 November 2022

Accepted 02 December 2022

Keywords:

Adherence

Medication

Treatment

Outcome

Myocardial infarction

© 2023 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Uso de polipíldora para mejorar la adherencia terapéutica y los resultados en adultos con infarto agudo de miocardio reciente: ¿es una estrategia efectiva?

INFO. ARTÍCULO
Historia del artículo:

Recibido 24 Octubre 2022

Recibido en forma revisada 19 Noviembre 2022

Aceptado 02 Diciembre 2022

Palabras clave:

Adherencia

Medicación

Tratamiento

Resultado

Infarto miocardio

© 2023 Los Autores. Publicado por Iberoamerican Journal of Medicine. Éste es un artículo en acceso abierto bajo licencia CC BY (<http://creativecommons.org/licenses/by/4.0/>).

HOW TO CITE THIS ARTICLE: Picón-Jaimes YA, Garcés-Lince MM, Gómez-Gómez JF, Villamizar-Rueda O. Use of polypill to improve therapeutic adherence and outcomes in adults with recent acute myocardial infarction: is it an effective strategy? Iberoam J Med. 2023;5(1):1-3. doi: 10.53986/ibjm.2023.0005.

Cardiovascular diseases remain the leading cause of morbidity and mortality in the world, generating an

* Corresponding author.

E-mail address: colmedsurg.center@gmail.com

ISSN: 2695-5075 / © 2023 The Authors. Published by Iberoamerican Journal of Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.53986/ibjm.2023.0005>

unsustainable global disease burden [1,2]. The World Health Organization estimates that approximately 18 million people die annually from cardiovascular complications (32% of all deaths worldwide, mainly from acute myocardial infarction and cerebrovascular disorders) and that more than 75% occur in low- and middle-income countries [1]. Because of this, the global health agenda defined as a priority the control of the most prevalent chronic non-communicable diseases, based on primary and secondary prevention strategies [2]. The polypill is a therapeutic strategy aimed at helping drug adherence, studied in depth in cardiovascular diseases [3, 4]. However, due to the social, cultural, economic and health differences in the different regions, it has not been precisely defined whether it is an effective and reproducible strategy, despite having shown promising results.

Very recently, Castellano et al [5] published the results of a randomized controlled trial, where they enrolled 2499 individuals with a history of acute myocardial infarction 6 months prior and administered the polypill strategy (Aspirin 100 mg, Ramipril 2.5/5/10 mg and Atorvastatin 20/40 mg) vs. usual care. Cardiovascular death, nonfatal acute myocardial infarction type 1, myocardial ischemia, or urgent need for revascularization were evaluated as outcomes. During a median follow-up of 36 months, 118/1237 (9.5%) major cardiovascular events occurred in the intervention group vs. 156/1229 (12.7%) in the control group (Hazard Ratio [HR] 0.76; 95% CI 0.60 - 0.96, $p=0.02$). The reported adherence was higher in the polypill group and adverse events were similar between the groups. This allowed the authors to conclude that the polypill was an effective and safe strategy in the prevention of major cardiovascular events in those with a recent history of acute myocardial infarction [5].

The same results have been obtained in other countries such as India [6] and Spain [3, 4], although the evidence supporting these studies is limited. Barrios et al [3] carried out a cost-effectiveness study in Spain on a simulated model, where they evaluated the 10-year cardiovascular outcomes in a male population with an average age of 64 years and a history of infarction, who would be administered polypill. They found that using this strategy would prevent 46 non-fatal events and 11 fatal events per 1000 individuals treated. Probability analysis showed that this strategy would be approximately 90% more effective than multiple monotherapies, saving thousands of euros per quality-adjusted life year [3]. In the same country, Ros-Castelló et al [4] evaluated the potential of the polypill on the secondary prevention of cerebrovascular disease in a cohort of 104 patients (54 received the polypill vs. 50 received conventional treatment). Their results showed a reduction in

systolic blood pressure in the intervention group ($p=0.002$), but without finding superiority in the effect on cholesterol (41% vs. 44%). Adverse events were moderate and similar in both groups, as well as adherence to treatment (93% vs. 88%) and stroke recurrence at 90 days (0% vs. 1%) [4]. This allowed the authors to conclude that there were no significant differences in adherence, as opposed to the reduction in systolic pressure. In India, Singh et al [6] conducted a cost-effectiveness study on the use of the polypill in secondary prevention vs. usual care, observing a significantly lower cost per patient with the polypill (-\$203, $p < 0.001$). According to variations, a price of \$0.94 per polypill was estimated, resulting to be cost-effective due to the outcomes obtained in previous studies [6].

In Latin America, particularly in Colombia [7] and Chile [8], studies have been carried out to measure drug adherence in secondary prevention for cardiovascular diseases, showing that it is less than 70% [7,8]. However, it is necessary to highlight that this adherence is affected by certain dimensions such as socioeconomic, provider-related, therapy experience-related and personal factors of the patient. Thus, 70% adherence cannot be expected in poor communities where out-of-pocket expenses do not allow for the independent cost of drugs, in areas where health systems fall short of providing drugs, where side effects and social myths prevent a rational weighing of the benefit-risk of therapy, and where scientific literacy and health education is scarce, which does not allow the patient to understand the relevance of adherence to treatment [7, 8].

Thus, and according to the current needs of global health that must be matched by evidence-based medicine [9-11], it is necessary to: 1) Design studies to obtain primary data on the behavior of this effect in each region, adapted to its social and health context; 2) To rigorously evaluate the methodological and reporting quality of the evidence on polypills in secondary prevention of cardiovascular disease; 3) Attempt to establish through public policies the use of the polypill, achieving an affordable purchase rate for health systems and providers, so that it can be reproducible and full coverage of the population in need can be obtained. This could be a tool with a substantial positive impact in the medium- and long-term on global cardiovascular health, especially in the regions with the greatest need for cardiometabolic health care.

1. CONFLICT OF INTERESTS

The authors have no conflict of interest to declare. The authors declared that this study has received no financial support.

2. REFERENCES

1. World Health Organization. *Cardiovascular Disease* [Internet]. Available in: https://www.who.int/health-topics/cardiovascular-diseases#tab=tab_1 (Accessed Sep 2022).
2. Heller O, Somerville C, Suggs LS, Lachat S, Piper J, Aya Pastrana N, et al. *The process of prioritization of non-communicable diseases in the global health policy arena. Health Policy Plan.* 2019;34(5):370-83. doi: 10.1093/heapol/cz043.
3. Barrios V, Kaskens L, Castellano JM, Cosin-Sales J, Ruiz JE, Zsolt I, et al. *Usefulness of a Cardiovascular Polypill in the Treatment of Secondary Prevention Patients in Spain: A Cost-effectiveness Study. Rev Esp Cardiol (Engl Ed).* 2017;70(1):42-9. doi: 10.1016/j.rec.2016.05.009.
4. Ros-Castelló V, Natera-Villalba E, Gómez-López A, Sánchez-Sánchez A, Chico-García JL, García-Madrona S, et al. *Use of the Cardiovascular Polypill in Secondary Prevention of Cerebrovascular Disease: A Real-Life Tertiary Hospital Cohort Study of 104 Patients. Cerebrovasc Dis Extra.* 2020;10(3):166-73. doi: 10.1159/000511064.
5. Castellano JM, Pocock SJ, Bhatt DL, Quesada AJ, Owen R, Fernandez-Ortiz A, et al. *Polypill Strategy in Secondary Cardiovascular Prevention. N Engl J Med.* 2022;387(11):967-77. doi: 10.1056/NEJMoa2208275.
6. Singh K, Crossan C, Laba TL, Roy A, Hayes A, Salam A, et al. *Cost-effectiveness of a fixed dose combination (polypill) in secondary prevention of cardiovascular diseases in India: Within-trial cost-effectiveness analysis of the UMPIRE trial. Int J Cardiol.* 2018;262:71-8. doi: 10.1016/j.ijcard.2018.03.082.
7. Ortega Oviedo SI, Vargas Rosero E. *Degree of adherence to treatments in people with cardiovascular risk. Av Enferm.* 2014; 32(1):25-32. doi: 10.15446/av.enferm.v32n1.46032.
8. Vejar M, Abufhele A, Varleta P, Araya MV, Escobar E, Fernández M, et al. *Adherence to pharmacologic treatment in secondary prevention of cardiovascular disease. Rev Chil Cardiol.* 2016; 35(3):270-82. doi: 10.4067/S0718-85602016000300010.
9. Mass-Hernández LM, Acevedo-Aguilar LM, Lozada-Martínez ID, Osorio-Agudelo LS, Maya-Betancourth JGEM, Paz-Echeverry OA, et al. *Undergraduate research in medicine: A summary of the evidence on problems, solutions and outcomes. Ann Med Surg (Lond).* 2022;74:103280. doi: 10.1016/j.amsu.2022.103280.
10. Lozada-Martínez ID, Suarez-Causado A, Solana-Tinoco JB. *Ethnicity, genetic variants, risk factors and cholelithiasis: The need for eco-epidemiological studies and genomic analysis in Latin American surgery. Int J Surg.* 2022;99:106589. doi: 10.1016/j.ijsu.2022.106589.
11. Pérez-Fontalvo NM, De Arco-Aragón MA, Jiménez-García JDC, Lozada-Martínez ID. *Molecular and computational research in low- and middle-income countries: Development is close at hand. J Taibah Univ Med Sci.* 2021;16(6):948-9. doi: 10.1016/j.jtumed.2021.06.010.